



Journal of Business & Economic Statistics

ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/ubes20

# **Dynamic Peer Groups of Arbitrage Characteristics**

Shuyi Ge, Shaoran Li & Oliver Linton

To cite this article: Shuyi Ge, Shaoran Li & Oliver Linton (2022): Dynamic Peer Groups of Arbitrage Characteristics, Journal of Business & Economic Statistics, DOI: 10.1080/07350015.2021.2011736

To link to this article: https://doi.org/10.1080/07350015.2021.2011736


View supplementary material



Published online: 28 Jan 2022.

٢	
L	

Submit your article to this journal 🖸





View related articles



View Crossmark data 🗹

Citing articles: 1 View citing articles 🕑

## **Dynamic Peer Groups of Arbitrage Characteristics**

Shuyi Ge<sup>a</sup>, Shaoran Li<sup>b</sup>, and Oliver Linton<sup>c</sup>

<sup>a</sup>School of Finance, Nankai University, Tianjin, China; <sup>b</sup>School of Economics, Peking University, Beijing, China; <sup>c</sup>Faculty of Economics, University of Cambridge, Cambridge, UK

#### ABSTRACT

We propose an asset pricing factor model constructed with semiparametric characteristics-based mispricing and factor loading functions. We approximate the unknown functions by B-splines sieve where the number of B-splines coefficients is diverging. We estimate this model and test the existence of the mispricing function by a power enhanced hypothesis test. The enhanced test solves the low power problem caused by diverging B-splines coefficients, with the strengthened power approaching one asymptotically. We also investigate the structure of mispricing components through Hierarchical K-means Clusterings. We apply our methodology to CRSP (Center for Research in Security Prices) and Compustat data for the U.S. stock market with one-year rolling windows during 1967–2017. This empirical study shows the presence of mispricing functions in certain time blocks. We also find that distinct clusters of the same characteristics lead to similar arbitrage returns, forming a "peer group" of arbitrage characteristics.

#### 1. Introduction

Stock returns have both common and firm-specific components. Ross (1976) proposed arbitrage pricing theory (APT) to summarize that expected returns on financial assets can be modeled as a linear combination of risk factors. In such a model, each asset has a sensitivity beta to the risk factor. The APT model explains the excess returns in the cross-sectional direction. Fama and French (1993) and Fama and French (2015) proxied those factors by the returns on portfolios sorted by different characteristics, and they developed three-factor and fivefactor models. After extracting the common movement parts, they treated the intercept as the mispricing alpha, which is assetspecific and cannot be explained by those risk factors. Many articles use a similar method to present other factor models, such as the four-factor model of Carhart (1997), the q-factor model of Hou et al. (2015), and the factor zoo by Feng et al. (2020) among others. All of the above articles studied observed factors and did not assign characteristics-based information to either alpha or beta.

Security-specific characteristics, such as capitalization and book to market ratio, are usually documented to explain assetspecific excess returns. Freyberger et al. (2020) analyzed the nonlinear effects of 62 characteristics through Lasso-style regressions. This study concluded that 13 of these characteristics have explanatory power on stock excess returns after selecting by adaptive group Lasso. Characteristics-based information is also exploited to develop arbitrage portfolios by directly parameterizing the portfolio weights as a linear function of characteristics, as in Hjalmarsson and Manchev (2012) and Kim et al. (2021). Empirically, they showed that their portfolio outperformed other baseline competitors.

This article's contributions are fourfold. First, we build up a more flexible semiparametric characteristics-based asset pricing factor model focusing on the mispricing component. Second, we extend previous estimation and testing methods, which can fit the current framework better. Especially, we extend the power-enhanced test of Fan et al. (2015) in a group manner to strengthen the conventional Wald test for mispricing functions. This test can also select the characteristics that contribute to arbitrage portfolios simultaneously. Third, we construct a twolayer clustering structure of mispricing components. Finally, our methods are applied to 50 years of monthly U.S. stock data. We detect distinct clusters of the same characteristics resulting in similar arbitrage returns, forming a "peer group" of arbitrage characteristics. This finding supplements existing portfolio management techniques by implying that the development of arbitrage portfolios through the asset weights determined by the linear mispricing function is improvable.

This class of models has a basic regression specification in Equation 1. Consider the panel regression model

$$y_{it} = \alpha_i + \sum_{j=1}^J \beta_{ji} f_{jt} + \epsilon_{it}, \qquad (1)$$

where  $y_{it}$  is the excess return of security *i* at time *t*;  $f_{jt}$  is the *j*th risk factor's return at time *t*;  $\beta_{ji}$  denotes the *j*th factor loading of asset *i*;  $\alpha_i$  represents the intercept (mispricing) of asset *i*; and  $\epsilon_{it}$  is the mean zero idiosyncratic shock. In terms of factor loadings

## ARTICLE HISTORY

Received July 2020 Accepted November 2021

KEYWORDS Characteristics-based; Peer groups; Power-enhanced test; Semiparametric

CONTACT Shaoran Li Shaoran@pku.edu.cn School of Economics, Peking University, Beijing 100871, China. Supplementary materials for this article are available online. Please go to www.tandfonline.com/UBES. 2022 American Statistical Association



 $\beta_{ji}$ , Connor and Linton (2007) and Connor et al. (2012) studied a characteristic-beta model, which bridges the beta-coefficients and firm-specific characteristics by specifying each beta as an unknown function of one characteristic. In their model, beta functions and unobservable factors are estimated by the backfitting iteration. They concluded that those characteristic-beta functions are significant and nonlinear. Their model can be summarized by

$$y_{it} = \sum_{j=1}^{J} g_j(X_{ji}) f_{jt} + \epsilon_{it}, \qquad (2)$$

where  $X_{ji}$  is the *j*th observable characteristic of firm *i*.

They restricted their beta function to be univariate and did not consider the components of factor loading functions that cannot be explained by characteristics. To overcome this limitation, Fan et al. (2016) allowed  $\beta_{ji}$  in Equation 2 to have a component explained by observable characteristics as well as an unexplained or stochastic part, written as  $\beta_{ji} = g_j(X_i) + u_{ji}$ , where  $u_{ji}$  is mean independent of  $X_i$ . They proposed the Projected Principal Component Analysis (PPCA), which projects stock excess returns onto space spanned by firm-specific characteristics and then applies Principal Component Analysis (PCA) to the projected returns to find the unobservable factors. This method has attractive properties even under the large *n* and small *T* setting. However, they did not study the mispricing part (alpha), which is crucial to both asset pricing theories and portfolio management.

In this article, we work on a semiparametric characteristicsbased alpha and beta model, which uses a set of security-specific characteristics that are similar to Freyberger et al. (2020). We use unknown multivariate characteristic functions to approximate both  $\alpha_i$  and  $\beta_{ji}$  in Equation (1). Specifically, we assume  $\alpha_i$  and  $\beta_{ji}$  are functions of a large set of asset-specific characteristics as  $\alpha_i = h(\mathbf{X}_i) + \gamma_i$  and  $\beta_{ji} = g_j(\mathbf{X}_i) + \lambda_{ij}^{-1}$ . We only specify additive structure of  $h(\mathbf{X}_i)$  and  $g_j(\mathbf{X}_i)$ , which are further approximated by B-splines sieve. We then estimate  $h(\mathbf{X}_i), g_j(\mathbf{X}_i)$  and unobservable risk factors  $f_{jt}$ . In addition, we design a power-enhanced test and Hierarchical K-mean Clustering for the mispricing function  $h(\mathbf{X}_i)$  to study the nonlinear behavior of arbitrage characteristics.

Some recent articles such as Kim et al. (2021) and Kelly et al. (2019) analyzed a similar model as ours, which assume that both  $h(X_i)$  and  $g_i(X_i)$  are *linear functions*. They both included around 40 characteristics in  $X_i$ . However, they drew different conclusions on the existence of  $h(X_i)$ . Kim et al. (2021) determined assets weights of arbitrage portfolios using oneyear rolling window estimates  $\frac{1}{n}\hat{h}(X_i)$ . They showed that their arbitrage portfolios returns are statistically and economically significant. However, Kelly et al. (2019) applied instrumented principal component analysis (IPCA) to the entire time span from 1965 to 2014, and concluded no evidence to reject the null hypothesis  $H_0$ :  $h(X_i) = X_i^{\mathsf{T}} B = 0$  through the bootstrapping. This dispute spurs the development of a more flexible model and reliable hypothesis tests to investigate the existence and structure of  $h(X_i)$ . The IPCA, which requires both large *n* and T to work, was introduced in Kelly et al. (2017). This method does not fit our setting since we apply rolling window analysis with small T. Furthermore, Kelly et al. (2019) restricted the function form of  $h(X_i)$  and  $g_i(X_i)$  to be time-invariant, which is not consistent with our empirical results under a semiparametric setting. To clarify the differences with the aforementioned research, this article proposes a semiparametric model, which allows for both nonlinearity and time-variation of  $h(X_i)$  and  $g_i(X_i)$ . Furthermore, we consider a different economic question, namely, the existence and structure of mispricing functions. Our empirical study sheds light on why Kelly et al. (2019) and Kim et al. (2021) drew different conclusions: weak, time-varying and nonlinear characteristics-based mispricing functions only appear in certain rolling windows, which is hard to be detected without rolling window analysis. However, given the presence of some persistent arbitrage characteristics, portfolios developed through mispricing functions can provide arbitrage returns.

The unrestrictive model in this article brings both opportunities and challenges. According to Huang et al. (2010), the number of B-spline knots must increase in the number of observations to achieve accurate approximation and good asymptotic performance. Therefore, the dimension of B-splines bases coefficients also needs to grow with the sample size. Besides, mispricing functions are treated as anomalies. Under a correctly specified factor model, coefficients of these B-splines bases that are employed to approximate  $h(X_i)$  are very likely to be sparse. All of these circumstances make the conventional Wald tests have very low power as discussed in Fan et al. (2015). Therefore, a power-enhanced test should be developed to strengthen the power of Wald tests and to detect the most relevant characteristics among a characteristic zoo included in  $h(X_i)$ . Kock and Preinerstorfer (2019) illustrated that if the number of coefficients diverges as the number of observations approaches infinity, the standard Wald test is power enhanceable. Fan et al. (2015) proposed a power-enhanced test by introducing a screening process on all estimated coefficients one by one. They added significant components as a supplement to the standard Wald test. In this article, we extend Fan et al. (2015) to a group manner to enhance the hypothesis test on a high-dimensional additive semiparametric function,  $H_0: h(X_i) = 0$ . This method allows all the significant components of  $h(X_i)$  to be selected and contribute to the test statistic, with the test power approaching one.

The careful analysis of  $h(X_i)$  is theoretically and practically meaningful. Firstly, the presence of  $h(X_i)$  is an important component of Arbitrage Pricing Theory (APT) and can contribute to asset pricing theories, namely, linking the mispricing functions with security-related characteristics. Secondly, as in Hjalmarsson and Manchev (2012) and Kim et al. (2021),  $h(X_i)$ can be used to construct arbitrage portfolios through observed characteristics. However, both research was built upon the condition that  $h(X_i)$  is linear over characteristics. If the mispricing function  $h(X_i)$  is not monotonic, simply setting portfolio weights to the estimated values of linear-specified  $h(X_i)$  can be problematic. In this article, we show that some characteristics with substantially different values give rise to similar arbitrage returns. The distance of arbitrage returns between two assets *i* and *j* is  $d_{ij} = |h(X_i) - h(X_j)|$  and the similarity of their characteristics is  $||X_i - X_i||_2$ , where  $|| \cdot ||_2$  represents  $L_2$  distance. Inspired by Hoberg and Phillips (2016) and Vogt and

 $<sup>{}^{1}</sup>X_{i}$  is a vector of a large set of asset-specific characteristics of stock *i*.

Linton (2017), we employ a hierarchical K-means clustering to classify these characteristics within each mispricing return group. We present the dynamic of distinct clusters of the same characteristics leading to similar arbitrage returns, forming a "peer group" of arbitrage characteristics. Therefore, under the semiparametric setting, the asset weighting function should rely on these time-varying and nonlinear peer groups.

The rest of this article is organized as follows. Section 2 sets out the semiparametric model. Section 3 introduces the assumptions and estimation methods. Section 4 constructs a powerenhanced test for high dimensional additive semiparametric functions. Section 5 employs Hierarchical K-Means Clustering to investigate peer groups of arbitrage characteristics. Section 6 describes the asymptotic properties of our estimates and test statistics. Section 6 simulates data to verify the performance of our methodology. Section 7 presents an empirical study. Finally, Section 8 concludes this article. Characteristics description tables, proofs, mispricing curves, and plots of peer groups are arranged in the appendix.

#### 2. Model Setup

We assume that there are n securities observed over T time periods. We also assume that during a short time window, each security has P time-invariant observed characteristics, such as market capitalization, momentum, and book-to-market ratios. Meanwhile, we may omit heteroscedasticity by assuming that each characteristic shares a certain form of variation within each period for all securities. We suppose that

$$y_{it} = (h(X_i) + \gamma_i) + \sum_{j=1}^{J} (g_j(X_i) + \lambda_{ij}) f_{jt} + \epsilon_{it}, \qquad (3)$$

where  $y_{it}$  is the monthly excess return of the *i*th stock at the month t;  $X_i$  is a 1  $\times$  P vector of P characteristics of stock i during time periods  $t = 1, \dots, T$ . T is a small and fixed time block. In practice, most characteristics are updated annually. Thus, we assume  $X_i$  is time-invariant in the one-year time window.  $h(X_i)$ is an unknown mispricing function explained by a large set of characteristics whereas  $\gamma_i$  is the random intercept of the mispricing part that cannot be explained by characteristics. Similarly, we have characteristics-beta function  $g_i(\cdot)$  to explain the *j*th factor loadings and the unexplained stochastic part of the loading is  $\lambda_{ij}$ .  $\lambda_{ij}$  is orthogonal to the  $g_i(\cdot)$  function.  $f_{jt}$  is the realization of the *j*th risk factor at time *t*. Finally,  $\epsilon_{it}$  is homoskedastic zeromean idiosyncratic residual of the *i*th stock at time *t*. Random variables  $\gamma_i$  and  $\lambda_{ii}$  are used to generalize our settings and not to be estimated. They will be treated as noise in the identification assumptions.

To avoid the curse of dimensionality, we impose additive forms on both  $h(\cdot)$  and  $g_j(\cdot)$  functions:  $h(\mathbf{X}_i) = \sum_{p=1}^{P} \mu_p(X_{ip})$ and  $g_j(\mathbf{X}_i) = \sum_{p=1}^{P} \theta_{jp}(X_{ip})$ , where  $\mu_p(X_{ip})$  and  $\theta_{jp}(X_{ip})$  are uni-variate unknown functions of the *p*th characteristic  $X_p$ . We rewrite the model as:

$$y_{it} = (\sum_{p=1}^{p} \mu_p(X_{ip}) + \gamma_i) + \sum_{j=1}^{J} (\sum_{p=1}^{p} \theta_{jp}(X_{ip}) + \lambda_{ij}) f_{jt} + \epsilon_{it},$$
(4)

Assumption 1. We suppose that

E

$$E(\epsilon_{it}|\mathbf{X}, f_{jt}) = 0,$$
  
$$(h(\mathbf{X}_i)) = E(g_j(\mathbf{X}_i)) = 0, \text{ for } j = 1, 2, \dots, J,$$
  
$$E(\gamma_i | \mathbf{X}) = E(\gamma_i),$$
  
$$E(\lambda_{ij} | \mathbf{X}) = E(\lambda_{ij}), \text{ for } j = 1, 2, \dots, J,$$
  
$$E(h(\mathbf{X}_i)g_j(\mathbf{X}_i)) = \mathbf{0}, \text{ for } j = 1, 2, \dots, J.$$

Similar to Connor et al. (2012) and Fan et al. (2016), Assumption 1 is to standardize model settings, including the zero mean assumption for factor loadings and mispricing functions for identification purposes. We also impose orthogonality between mispricing and factor loading parts for the identification reason. This is because the variation of risk factors can be absorbed into the mispricing part if it is not orthogonal to the factor loadings. More discussions can be found in Connor et al. (2012).

#### 3. Estimation

In this section, we discuss the approximation of unknown univariate functions and our estimation methods for model Equation 3. In the semiparametric setting, we apply the Projected-PCA following Fan et al. (2016) to work on the common factors and characteristics-beta directly. Next, we project the residuals onto the characteristics-based alpha space that is orthogonal to the systematic part. The second step is similar to equalityconstrained OLS.

#### 3.1. B-splines Approximation

We use B-splines sieve to approximate unknown functions  $\theta(\cdot)$ and  $\mu(\cdot)$  in Equation (4). Similar to Huang et al. (2010) and Chen and Pouzo (2012), we have the following procedures. First, suppose that the *p*th covariate  $X_p$  is in the interval  $[D_0, D]$ , where  $D_0$  and D are finite numbers with  $D_0 < D$ . Let  $\mathbf{D} = \{\underbrace{D_0, D_0, \ldots, D_0}_{l} < d_1 < d_2 < \cdots < d_{m_n} < \underbrace{D, D, \ldots, D}_{l}\}$  be a simple knot sequence on the interval  $[D_0, D]$ . Here,  $m_n = \lfloor n^{\nu} \rceil$  $(\lfloor \cdot \rceil$  gives nearest integer) is a positive integer of the number of internal knots, which is a function of security size *n* in period *t* with  $0 < \nu < 0.5$ . *l* is the degree of those bases. Therefore, we have  $H_n = l + m_n$  bases in total, which will diverge as  $n \to \infty$ . Following this setting, a set of B-splines can be built for the space  $\Omega_n[\mathbf{D}]$ .

Second, for the *p*th characteristic  $X_p$ , there is a set of  $H_n$  orthogonal bases  $\{\phi_{1p}(X_p), \ldots, \phi_{H_np}(X_p)\}$ . Those univariate unknown functions can be approximated as linear combinations of these bases as  $\mu_p(X_p) = \sum_{q=1}^{H_n} \alpha_q \phi_{qp}(X_p) + R_p^{\mu}(X_p)$  and  $\theta_p(X_p) = \sum_{q=1}^{H_n} \beta_{jq} \phi_{qp}(X_p) + R_p^{\theta}(X_p)$ , where  $R_p^{\mu}(X_p)$  and  $R_p^{\theta}(X_p)$  are approximation errors. It is not necessary to use the same bases for both unknown functions and the

representation here is for notational simplicity only. Therefore, the model Equation (4) can be written as follows:

$$y_{it} = \sum_{p=1}^{P} (\sum_{q=1}^{H_n} \alpha_{pq} \phi_{pq}(X_{ip}) + R_p^{\mu}(X_p)) + \gamma_i + \sum_{j=1}^{J} (\sum_{p=1}^{P} (\sum_{q=1}^{H_n} \beta_{jpq} \phi_{pq}(X_{ip}) + R_p^{\theta}(X_p)) + \lambda_{ij}) f_{jt} + \epsilon_{it}.$$

For each i = 1, 2, ..., n, p = 1, 2, ..., P and t = 1, 2, ..., T, we have

$$\mathbf{1}_T = (1, \ldots, 1)^\mathsf{T} \in \mathbb{R}^T,$$

$$\beta_j = (\beta_{1,j1}, \dots, \beta_{H_n,j1}, \dots, \beta_{1,jP}, \dots, \beta_{H_n,jP})^\mathsf{T} \in \mathbb{R}^{H_n P},$$

$$\boldsymbol{B}=(\beta_1,\ldots,\beta_J),$$

$$\mathbf{A} = (\alpha_{11}, \ldots, \alpha_{1H_n}, \ldots, \alpha_{P1}, \ldots, \alpha_{PH_n})^{\mathsf{T}} \in \mathbb{R}^{H_n P},$$

$$\Phi(\mathbf{X}) = \begin{bmatrix} \phi_{1,11}(X_{11}) \cdots \phi_{1,1H_n}(X_{11}) \cdots \phi_{1,P1}(X_{1P}) \cdots \phi_{1,PH_n}(X_{1P}) \\ \phi_{2,11}(X_{21}) \cdots \phi_{2,1H_n}(X_{21}) \cdots \phi_{2,P1}(X_{2P}) \cdots \phi_{2,PH_n}(X_{2P}) \\ \vdots & \vdots & \ddots & \vdots \\ \phi_{n,11}(X_{n1}) \cdots \phi_{n,1H_n}(X_{n1}) \cdots \phi_{n,P1}(X_{nP}) \cdots \phi_{n,PH_n}(X_{nP}) \end{bmatrix},$$

where  $\phi_{i,ph}(X_{ip})$  is the *h*th basis of the *p*th characteristic of asset *i* at time *t*. Therefore, the original model

$$Y = (h(X) + \Gamma)\mathbf{1}_T^{\mathsf{T}} + (G(X) + \Lambda)F^{\mathsf{T}} + U,$$

can be represented by B-splines sieve as follows:

$$Y = (\Phi(X)A + \Gamma + R^{\mu}(X))\mathbf{1}_{T}^{\mathsf{T}} + (\Phi(X)B + \Lambda + R^{\theta}(X))F^{\mathsf{T}} + U.$$
(5)

*Y* is the  $n \times T$  matrix of  $y_{it}$ ;  $\Phi(X)$  is the  $n \times PH_n$  matrix of Bsplines bases; *A* is a  $PH_n \times 1$  matrix of mispricing coefficients;  $\mathbb{R}^{\mu}(X)$  is an  $n \times 1$  matrix of approximation errors; *B* is a  $PH_n \times J$ matrix factor loadings' coefficients;  $\mathbb{R}^{\theta}(X)$  is a  $n \times J$  matrix of approximation errors. We have  $\mathbb{R}_p^{\mu}(X_p) \rightarrow^p 0$  and  $\mathbb{R}_p^{\theta}(X_p) \rightarrow^p$ 0, as  $n \to \infty$  as in Huang et al. (2010). Therefore, we omit the approximation errors for simplicity henceforth. *F* is the  $T \times J$ matrix of  $f_{tj}$  and *U* is a  $n \times T$  matrix of  $\epsilon_{it}$ . h(X) is a  $n \times 1$  vector of characteristics-based mispricing component; G(X) is an  $n \times J$ vector of characteristics-based factor loadings;  $\mathbf{1}_T$  is a  $T \times 1$ vector of 1. The rest are defined the same as Equation 4.

We define a projection matrix as follows:

$$\boldsymbol{P} = \boldsymbol{\Phi}(\boldsymbol{X})(\boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}}\boldsymbol{\Phi}(\boldsymbol{X}))^{-1}\boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}}.$$

The remaining goals of this article are to estimate both h(X) and G(X) consistently and conduct a power-enhanced test on the hypothesis  $H_0$ : h(X) = 0, i.e., to check the existence of mispricing functions under semiparametric settings. Finally, we cluster peer groups of arbitrage characteristics.

#### 3.2. Two-Step Projected-PCA

In this section, we combine and extend Projected-PCA by Fan et al. (2016) and equality constrained least square similar to Kim et al. (2021) to estimate the model. To facilitate the estimation, we define a  $T \times T$  time series demeaning matrix  $D_T = I_T - \frac{1}{T} \mathbf{1}_T \mathbf{1}_T^T$ .<sup>2</sup> Next, we demean the equation above on both sides. Therefore, we have

$$YD_T = \tilde{Y} = (\Phi(X)B + \Lambda)F^{\mathsf{T}}D_T + UD_T.$$

Mispricing terms disappear since they are time-invariant by  $(\Phi(X)A + \Gamma)\mathbf{1}_T^{\mathsf{T}}D_T = \mathbf{0}$ . This helps us to work on the systematic part later. Henceforth, we use F to represent the time-demeaned factor matrix.

Our procedures are designed to estimate factor loadings G(X), time-demeaned unobserved factors F and mispricing coefficients A in sequence.

Under Assumption 1, we have the following estimation procedures:

- 1 Projecting  $\tilde{Y}$  onto the spline space spanned by  $\{X_{ip}\}_{i \leq n, p \leq P}$ through a  $n \times n$  projection matrix P with  $P = \Phi(X)(\Phi(X)^{\mathsf{T}} \Phi(X))^{-1}\Phi(X)^{\mathsf{T}}$ . We then collect the projected data  $\hat{Y} = \Phi(X)(\Phi(X)^{\mathsf{T}}\Phi(X))^{-1}\Phi(X)^{\mathsf{T}}\tilde{Y}$ .
- 2 Applying the principal component analysis to the projected data  $\hat{Y}^{\mathsf{T}}\hat{Y}$ . This allows us to work directly on the sample covariance of  $G(X)F^{\mathsf{T}}$ , under the condition  $E(g_j(X_i)\epsilon_{it}) = E(g_j(X_i)\lambda_{ij}) = 0$ .
- 3 Estimating  $\hat{F}$  as the eigenvectors corresponding to the first *J* (assumed given) largest eigenvalues of the  $T \times T$  matrix  $\frac{1}{n} \hat{Y}^{\mathsf{T}} \hat{Y}$  (covariance of projected  $\hat{Y}$ ).

The method above substantially improves estimation accuracy and facilitates theoretical analysis even under the large n and small T. Small T is preferable in our model setting as we use one-year rolling windows analysis in both simulation and empirical studies, and large n is required for asymptotic analysis.

Factor loadings  $\hat{G}(X)$  are estimated as follows:

$$\hat{G}(X) = \hat{Y}\hat{F}(\hat{F}^{\mathsf{T}}\hat{F})^{-1}$$

In the next step, we estimate the coefficients of the mispricing bases.

4 The estimator of *A* is

$$\hat{A} = \arg\min_{A} \operatorname{vec}(Y - \Phi(\mathbf{X})A\mathbf{1}_{T}^{\mathsf{T}} - \hat{G}(\mathbf{X})\hat{F}^{\mathsf{T}})^{\mathsf{T}}$$
$$\operatorname{vec}(Y - \Phi(\mathbf{X})A\mathbf{1}_{T}^{\mathsf{T}} - \hat{G}(\mathbf{X})\hat{F}^{\mathsf{T}}), \qquad (6)$$

subject to  $\hat{G}(X)^{\mathsf{T}} \Phi(X) A = \mathbf{0}_{\mathsf{J}}$ .

Let a  $PH_n \times 1$  vector  $\hat{A}$  be a closed-form solution:

 $\hat{A} = Q\tilde{A},$ 

where

$$Q = I - (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1} \Phi(X)^{\mathsf{T}} \hat{G}(X) (\hat{G}(X)^{\mathsf{T}} \hat{G}(X))^{-1} \hat{G}(X)^{\mathsf{T}} \Phi(X),$$

<sup>2</sup> $I_T$  is a  $T \times T$  identity matrix, and  $\mathbf{1}_T$  is a  $T \times 1$  matrix of 1.

$$\tilde{A} = \frac{1}{T} (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1} \Phi(X)^{\mathsf{T}} (Y - \hat{G}(X)\hat{F}^{\mathsf{T}}) \mathbf{1}_{T},$$

given  $P\hat{G}(X) = \hat{G}(X)$ .

As in Assumption 1, the h(X) is orthogonal to the characteristics-based loadings G(X).

5 We also estimate the covariance matrix of  $\hat{A}$ , that is,  $\Sigma$ , by extending the methods of Liew (1976). This can facilitate theoretical analysis in the next section. According to Liew (1976),  $\hat{A}$  is the equality constrained least-square estimator, which has the covariance matrix as (only under  $n \leq T$  and covariance shrinkage as in Ledoit et al. (2012) and Fan et al. (2013) among others.):

$$\mathbf{\hat{P}} = \mathbf{Q}\mathbf{\hat{P}}_{\tilde{A}}\mathbf{Q}^{\mathsf{T}},$$

where:

$$\mathfrak{D}_{\tilde{A}} = (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1} \Phi(X)^{\mathsf{T}} \begin{bmatrix} \hat{\sigma}_{1}^{2} & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & \hat{\sigma}_{n}^{2} \end{bmatrix}$$

$$\Phi(X) (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1},$$

$$\hat{\sigma}_i^2 = \frac{\sum_{1}^{T} \hat{e}_{it}^2}{T}$$

where  $\sum_{1}^{T} \hat{e}_{it}^2 = \sum_{1}^{T} (y_{it} - \bar{y}_i - \sum_{p=1}^{P} \sum_{q=1}^{H_n} \hat{\alpha}_{pq} \phi_{pq}(x_{ip}) - \sum_{j=1}^{J} (\sum_{p=1}^{P} \sum_{q=1}^{H_n} \hat{\beta}_{jpq} \phi_{pq}(x_{ip})) \hat{f}_{jt})^2$ . Heteroscedasticity is caused by  $\gamma_i$ .

#### 4. Power-Enhanced Tests

There are considerable discussions about the mispricing phenomenon under factor models, while the existence of mispricing functions remains controversial. Namely, whether there are relevant covariates explaining remaining excess returns after subtracting co-movements components captured by risk factors. Recently, Kim et al. (2021) found the characteristics arbitrage opportunities by estimating a linear characteristic mispricing function without providing theoretical results. However, Kelly et al. (2019) conducted a conventional Wald hypothesis test on the similar mispricing function using bootstrap, concluding that there is no evidence to reject the null hypothesis  $H_0$ : h(X) =**0**. Additionally, they applied the bootstrap method to estimate the covariance matrix  $\Sigma$ , which caused potential problems for theoretical analysis. Moreover, according to Fan et al. (2015), their test results may have relatively low power when the true coefficient vector of linear mispricing function A has a sparse structure.

Both studies adopt a parametric framework, which relies on the strong assumption of linearity. However, this assumption is not consistent with Connor et al. (2012), which showed that both characteristic-beta and mispricing functions are very likely to be nonlinear. Therefore, we propose a semiparametric model to accommodate the nonlinearity to a great extent.

But semiparametric framework leads to additional challenges for inference. On the one hand, as mentioned above, the number of coefficients of mispricing B-splines diverges as  $n \rightarrow \infty$ , which implies that the power of the standard Wald test can be quite low (see Fan et al. 2015). On the other hand, according to other research like Fama and French (1993) and Fama and French (2015), mispricing terms can be regarded as anomalies. This means that in our model setting, the true mispricing coefficient vector A can be high-dimensional but sparse, reducing the power of the conventional Wald test further.

As in Kock and Preinerstorfer (2019), conventional hypothesis tests under these circumstances are power enhanceable. The power-enhanced Wald test in this article is an extension of Fan et al. (2015) to a group manner, namely, the hypothesis test under high-dimensional additive semiparametric settings. The proposed test is power strengthened when the dimension of the coefficients of the additive regression A is diverging as  $n \to \infty$ without size distortion. Meanwhile, this test is robust to sparse alternatives. On top of that, the proposed test can select the most important components from sparse additive functions. Finally, the proposed method can also be applied when the number of characteristics is diverging, that is,  $P \to \infty$ .

We construct a new test

$$H_0: h(\mathbf{X}) = \mathbf{0}, \qquad H_1: h(\mathbf{X}) \neq \mathbf{0},$$

equivalently,

$$H_0: \boldsymbol{A} = \boldsymbol{0}, \qquad H_1: \boldsymbol{A} \in \mathcal{A},$$

where  $\mathcal{A} \subset \mathbb{R}^{PH_n} \setminus \mathbf{0}$ . Here, we have:

$$S_1 = \frac{\hat{A}^{\mathsf{T}} \mathbf{\hat{Q}}^{-1} \hat{A} - PH_n}{\sqrt{2PH_n}},$$

where  $S_1$  is the "original" Wald test statistic; P is the number of characteristics;  $PH_n$  is the total number of B-spline bases, and  $\hat{A} \in \mathbb{R}^{PH_n}$ . The value of  $H_n$  is a function of asset number n, and  $H_n \to \infty$  as  $n \to \infty$ . Under  $H_0$ ,  $S_1$  has a nondegenerate limiting distribution F as  $n \to \infty$ . Given the significance level  $q, q \in (0, 1)$  as well as the critical value  $F_q$ :

$$S_1|H_0 \to^d F,$$
  
$$\lim_{n \to \infty} \Pr(S_1 > F_q|H_0) = q.$$

Pesaran and Yamagata (2012) showed that

$$S_1|H_0 \rightarrow^a \mathcal{N}(0,1),$$

under regularity conditions.

Potentially, sparse and diverging  $PH_n$  means that it is plausible to add a power-enhanced component to  $S_1$ , which can improve the power of the hypothesis test without any size distortions.

Therefore, we can construct an extra screening component  $S_0$  as follows:

$$S_0 = H_n \sum_{p=1}^{P} I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n),$$

where  $\hat{\sigma}_{ph}$  is the square-root of the *ph*th entry of the diagonal elements of  $\hat{\Sigma}$ .  $I(\cdot)$  is an indicator for the screening process while  $\eta_n$  is a data-driven threshold value to avoid potential size-distortion.

Here, we discuss the choice of  $\eta_n$ . By construction and the assumption of independent characteristics, we assume that all B-splines bases are orthogonal. Our goal is to bound the maximum of those standardized coefficients under the null hypothesis.

Define  $Z = \max_{\{1 \le p \le P, 1 \le h \le H_n\}} \{|\hat{\alpha}_{ph}| / \hat{\sigma}_{ph}\}$ . We have

$$\hat{\alpha}_{ph}/\hat{\sigma}_{ph}|H_0 \rightarrow^d N(0,1),$$
  
 $E(Z|H_0) = \sqrt{2\log PH_n}.$ 

After grouping the coefficients of bases used to approximate the unknown function of each characteristic, we try to bound  $\max(\sum_{h=1}^{H_n} |\hat{\alpha}_{1h}| / \hat{\sigma}_{1h}, \dots, \sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph}, \dots, \sum_{h=1}^{H_n} |\hat{\alpha}_{Ph}| / \hat{\sigma}_{Ph})$ under  $H_0$ , Following this, we set the threshold as  $\eta_n = H_n \sqrt{2 \log(PH_n)}$ , where  $H_n = l + n^{\nu}$ . As  $H_n$  is a diverging sequence, it can control the influence of the group size properly. Meanwhile,  $\eta_n$  also diverges so that  $\eta_n$  is a conservative threshold value used to avoid potential size distortion.

Apart from strengthening the power of conventional hypothesis test,  $I(\cdot)$  is a screening term that can select the most relevant characteristics at the same time.

We then define the arbitrage characteristics set, which includes the characteristics that have the strong explanation power for mispricing functions:

$$\hat{\mathcal{M}} = \{ \boldsymbol{X}_{\boldsymbol{p}} \in \boldsymbol{X} : \sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n, \quad \boldsymbol{p} = 1, 2, \dots, P \},\$$

and *M* is the cardinality of the set containing mispricing characteristics. When the set  $\hat{\mathcal{M}}$  is relatively small, conventional tests are likely to suffer the lower power problem. The added  $S_0$  strengthens the power of the test and drives the power to one since  $S_0$  is slowly diverging.

Therefore, our new test statistic is  $S = S_0 + S_1$ , and asymptotic properties of *S* will be discussed later.

To conclude, the advantages of  $S = S_0 + S_1$  are:

- 1. The power of the hypothesis test on  $H_0: h(X) = 0$  is mainly enhanced without size distortions.
- 2. We can find specific characteristics which cause the mispricing by this screening mechanism.

As designed,  $S_0$  satisfies all three properties of Fan et al. (2015), as  $n \to \infty$ :

- 1.  $S_0$  is nonnegative,  $Pr(S_0 \ge 0) = 1$
- 2.  $S_0$  does not cause size distortion: under  $H_0$ ,  $Pr(S_0 = 0|H_0) \rightarrow 1$
- 3.  $S_0$  enhances test power. Under  $H_1$ ,  $S_0$  diverges quickly in probability given the well chosen  $\eta_n$ .

Based on properties of  $S_0$ , we have three properties of S listed

1. No size-distortion  $\limsup_{n \to \infty} \Pr(S > F_q | H_0) = q$ 

- 2.  $Pr(S > F_q|H_1) \ge Pr(S_1 > F_q|H_1)$ . Hence, the power of *S* is at least as large as that of *S*<sub>1</sub>.
- 3.  $\Pr(S > F_q | \hat{\mathcal{M}} \neq \emptyset) \rightarrow 1$  when  $S_0$  diverges. This happens, especially, when the true form of *A* has a sparse structure.

#### 5. Hierarchical K-Means Clustering

This section introduces a Hierarchical K-means Clustering method to find peer groups of arbitrage characteristics based on their arbitrage returns. We ask whether distinct groups of the same arbitrage characteristics, according to their similarity measured by  $||X_i - X_j||_2$ , may result in similar characteristicbased arbitrage returns in each rolling block, which is an implication for the nonmonotonic mispricing function, and forms a "peer group" of arbitrage characteristics. Because traditional arbitrage portfolios as in Kim et al. (2021) and Hjalmarsson and Manchev (2012) relied on the linearity of characteristics-based mispricing components to work, our clustering results can show whether this method is still applicable under a more flexible semiparametric model. If there are persistent peer groups in arbitrage returns, investors should consider to long the assets in the peer group with the highest arbitrage returns while short the assets in the peer group with the lowest arbitrage returns to form an arbitrage portfolio.

Introduction of K-means clustering can be found in Cox (1957) and Fisher (1958).

After the screening process in Section 4, we obtain the relevant components of mispricing function h(X), which is estimated as

$$\hat{\mathcal{M}} = \{ \boldsymbol{X}_{\boldsymbol{p}} \in \boldsymbol{X} : \sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n, \quad p = 1, 2, \dots, P \}.$$

We define an  $n \times M$  matrix M of arbitrage characteristics at time window t as follows:

$$M = \{X_1, \ldots, X_m, \ldots, X_M\}, \text{ where } X_m \in \hat{\mathcal{M}}.$$

Note that these characteristics are time-invariant in each rolling window. We also set characteristics-based arbitrage returns of asset i in month t as:

$$\ddot{y}_{it} = \boldsymbol{\phi}(\boldsymbol{M}_i) \hat{A}_M,$$

where  $\phi(M_i)$  and  $\hat{A}_M$  are the corresponding parts of the matrix  $\Phi(X_i)$  and the vector  $\hat{A}$ . For each rolling window, we classify all n assets through a 2-layer K-means clustering. At the first layer, we cluster these assets into K groups according to the similarity of their characteristics-based arbitrage returns  $\ddot{y}_{it}$ . At the second layer, we divide  $R_j$  subgroups within the jth group from the first layer by the similarity of their arbitrage characteristics, where  $j = 1, 2, \ldots, K$ . Finally, the peer groups of arbitrage characteristics can be attained. We repeat this method for all rolling blocks to investigate dynamic patterns of these peer groups. These clusterings will provide illustrative evidence of linear/nonlinear and time-invariant/time-varying structure of mispricing function h(X).

We give the classification procedures of both layers. We define  $\Delta_{ij}$  as the difference between characteristics-based arbitrage returns of  $\ddot{y}_{it}$  and  $\ddot{y}_{jt}$ , as well as  $\Upsilon_{ij}$  as the difference between arbitrage characteristics

$$\Delta_{ij} = \ddot{y}_{it} - \ddot{y}_{jt}$$
, where  $i \neq j, i, j = 1, 2, ..., n$ .  
 $\Upsilon_{ij} = ||M_i - M_j||_2$ , where  $i \neq j, i, j = 1, 2, ..., n$ ,

 $M_i$  represents the *i*th row of M. We set two tolerance thresholds  $\psi_y$  and  $\psi_x$ , which are used to control the biggest difference within each group of both layers separately. To accelerate the convergence of the K-means Clustering, we firstly apply a first difference process, which is introduced below, to obtain centroids as in Vogt and Linton (2017).

For the first layer, we have the first difference process:

- 1. *First difference*: We randomly pick *i*th asset and then we calculate  $\Delta_{ij}$  with other assets j = 1, 2, ..., n. Thus, we obtain  $\Delta_{i(1)} ... \Delta_{i(n)}$ , with *n* being the total individuals for classification. Without loss of generality, we assume  $\Delta_{i(1)} = \min{\{\Delta_{i(1)}, ..., \Delta_{i(n)}\}}$ , and  $\Delta_{i(n)} = \max{\{\Delta_{i(1)}, ..., \Delta_{i(n)}\}}$ .
- 2. Ordering: We rank the values obtained in Step 1 as follows:

$$\Delta_{i(1)} \leqslant \dots \leqslant \Delta_{i(j_1-1)} < \Delta_{i(j_1)} \leqslant \dots \leqslant \Delta_{i(j_2-1)}$$
$$< \Delta_{i(j_2)} \leqslant \dots \leqslant \Delta_{i(j_3-1)}$$
$$\vdots$$
$$< \Delta_{i(i_{k'-1})} \leqslant \dots \leqslant \Delta_{i(m)}.$$

We use the strict inequality mark to show large jumps of "first difference," all of which are larger than  $\psi_y$ , while the weak inequality means that the distance calculated is not larger than  $\psi_y$ . We identify K - 1 jumps that are larger than  $\psi_y$  above. Thus, the initial classification is achieved, and we have a total of K groups with  $j_1 - 1$  members in the first group  $C_1$ ,  $j_2 - j_1$  members in the second group  $C_2$ , ..., and  $n - j_{K-1} + 1$  members in the final group  $C_K$ .

In terms of the second layer, for the assets in the *k*th group  $C_k$ , we use the same method to further divide them into *r* 

subgroups as  $\mathcal{R}_{1k}, \mathcal{R}_{2k}, \ldots, \mathcal{R}_{rk}$ . Within each subgroup, we have

$$\Upsilon_{ab} = ||\mathbf{M}_a - \mathbf{M}_b||_2 \leq \psi_x, \text{ where } a, b \in \mathcal{R}_{ik}, i = 1, 2, \dots, r,$$
  
and  $k = 1, 2, \dots, K.$ 

The K-means algorithm is:

- 1. Step 1: Determine the starting mean values for each group  $\hat{c}_1^{[0]}, \ldots, \hat{c}_K^{[0]}$  and calculate the distances  $\hat{D}_k(i) = \Delta(\ddot{y}_{it}, \hat{c}_k^{[0]}) = |\ddot{y}_{it} \hat{c}_k^{[0]}|$  for each *i* and *k*. Define the partition  $\{C_1^{[0]}, \ldots, C_K^{[0]}\}$  by assigning the *i*th individual to the *k*-th group  $C_k^{[0]}$  if  $\hat{D}_k(i) = \min_{1 \le k' \le K} \hat{D}_{k'}(i)$ .
- 2. Step *l*: Let  $\{C_1^{[l-1]}, \ldots, C_K^{[l-1]}\}$  be the partition of  $\{1, \ldots, n\}$  from the latest iteration step. Calculate mean functions

$$\hat{\overline{c}}_k^{[l]} = \frac{1}{|\mathcal{C}_k^{[l-1]}|} \sum_{i \in \mathcal{C}_k^{[l-1]}} \ddot{y}_{it} \text{ for } 1 \leq k \leq K.$$

And then we calculate  $\Delta(\ddot{y}_{it}, \hat{c}_k^{[l]}) = |\ddot{y}_{it} - \hat{c}_k^{[l]}|$  for each *i* and *k*. Define the partition  $\{\mathcal{C}_1^{[l]}, \ldots, \mathcal{C}_K^{[l]}\}$  by assigning the *i*th individual to the *k*th group  $\mathcal{C}_k^{[l]}$  if  $\hat{D}_k(i) = \min_{1 \le k' \le K} \hat{D}_{k'}(i)$ .

3. Iterate the above steps until the partition  $\{C_1^{[w]}, \ldots, C_K^{[w]}\}$  does not change anymore.


To accelerate the convergence of K-means algorithm, at the step 1, results of the first difference are used. As we have already obtained our initial grouping  $\{C_1, \ldots, C_K\}$ , therefore, the starting value for the Step 1 is

$$\hat{\bar{c}}_{k}^{[0]} = \frac{1}{|\mathcal{C}_{k}|} \sum_{i \in \mathcal{C}_{k}} \ddot{y}_{it} \text{ for } 1 \leq k \leq K,$$

where  $|C_k|$  is the cardinality of the group  $C_k$ .

The consistency and other theoretical results of the above procedures can be found in Pollard (1981), Pollard et al. (1982), Sun et al. (2012) and Vogt and Linton (2017).

For the second layer, we repeat the procedures within each group  $C_k^{[w]}$  with respect to  $\Upsilon_{ab}$ , and the structure of characteristics-based arbitrage returns is:



The first layer is the structure of characteristics-based arbitrage returns, while the second layer gives peer groups of characteristics that can provide similar characteristics-based arbitrage returns.

The number of clusterings is determined by threshold values  $\psi_y$  and  $\psi_x$  directly.  $\psi_y$  and  $\psi_x$  are chosen by the tradeoff between the number of clusterings and the total within-group sum of squares.

#### 6. Asymptotic Properties

This section discusses assumptions and properties of estimates and the power enhanced statistic *S*.

*Definition 6.1.* We define  $A \to^{p} B$  as  $n \to \infty$  of two  $n \times m$  matrix A and B with fixed p when  $\frac{1}{n}(A-B)^{\mathsf{T}}(A-B) \to^{p} \mathbf{0}_{m \times m}$  as  $n \to \infty$ .

#### 6.1. Assumptions

*Assumption 2.* As  $n \to \infty$ , we have

$$\frac{1}{n} Y^{\mathsf{T}} Y \to_P M_Y,$$
$$F^{\mathsf{T}} F = I_I,$$

where  $M_Y$  is a positive definite matrix, and  $I_J$  is a  $J \times J$  identity matrix.

We define  $\lambda_{min}(M)$  and  $\lambda_{max}(M)$  as the largest and the smallest eigenvalues of matrix M, respectively. Additionally, we define  $C_{min}$  and  $C_{max}$  to be positive constants such that

$$C_{min} \leq \lambda_{min}(\frac{1}{n}\Phi^{\mathsf{T}}(\mathbf{X})\Phi(\mathbf{X})) < \lambda_{max}(\frac{1}{n}\Phi^{\mathsf{T}}(\mathbf{X})\Phi(\mathbf{X})) \leq C_{max}$$
  
as  $n \to \infty$ .

We impose these restrictions to avoid non-invertibility of stock returns, characteristics, and rotation indeterminacy.

#### Assumption 3.

$$\frac{1}{n} \boldsymbol{G}(\boldsymbol{X})^{\mathsf{T}} \boldsymbol{G}(\boldsymbol{X}) \to_{P} \begin{bmatrix} d_{1} & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & d_{J} \end{bmatrix},$$

as  $n \to \infty$ , where  $d_j$  are distinct and positive entries.

Both Assumption 2 and 3 are similar to those in Fan et al. (2016), which are used to separately identify risk factors and factor loadings. Given the orthogonal bases of B-splines and uncorrelated or weakly correlated characteristics, Assumption 3 is mild.

Assumption 4.  $K_{\min}$  and  $K_{\max}$  are positive constants such that:

$$K_{\min} \leq \lambda_{\min}(\frac{1}{n}G(X)^{\mathsf{T}}PG(X)) < \lambda_{\max}(\frac{1}{n}G(X)^{\mathsf{T}}PG(X)) \leq K_{\max}$$
  
as  $n \to \infty$ .

This assumption requires nonvanishing explanatory power of the B-splines bases  $\Phi(X)$  on the factor loading matrix G(X).

Assumption 5.  $\epsilon_{it}$  is realized iid idiosyncratic shocks with  $E(\epsilon_{it}) = 0$  and  $var(\epsilon_{it}) = \sigma^2$ .

Heteroscedasticity is caused by  $\gamma_i$ , namely,  $var(\gamma_i + \epsilon_{it}) = \sigma_i^2$ .

#### 6.2. Main Results

*Theorem 6.1.* Let  $\hat{F}$  be the  $T \times J$  matrix estimate of latent risk factors. Under Assumption 1–4,  $\hat{F} \rightarrow^{P} F$ , as  $n \rightarrow \infty$ .

*Theorem 6.2.* Define the  $n \times J$  matrix  $\hat{G}(X)$  as the estimate of factor loadings G(X). Under Assumption 1–4 and Theorem 6.2, as  $n \to \infty$ , then  $\hat{G}(X) \to {}^{P} G(X)$ .

*Theorem 6.3.* Let the  $PH_n \times 1$  vector  $\hat{A}$  be the solution of constrained OLS Equation (6), then

$$\hat{A} = Q\tilde{A}$$

where

$$\boldsymbol{Q} = \boldsymbol{I} - (\boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}} \boldsymbol{\Phi}(\boldsymbol{X}))^{-1} \boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}} \hat{\boldsymbol{G}}(\boldsymbol{X}) (\hat{\boldsymbol{G}}(\boldsymbol{X})^{\mathsf{T}} \hat{\boldsymbol{G}}(\boldsymbol{X}))^{-1} \hat{\boldsymbol{G}}(\boldsymbol{X})^{\mathsf{T}} \boldsymbol{\Phi}(\boldsymbol{X}),$$

$$\tilde{A} = \frac{1}{T} (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1} \Phi(X)^{\mathsf{T}} (Y - \hat{G}(X)\hat{F}^{\mathsf{T}}) \mathbf{1}_{T}.$$

Under Assumption 1–4,  $\Phi(\mathbf{X})\hat{\mathbf{A}} \rightarrow^{P} h(\mathbf{X})$ , as  $n \rightarrow \infty$ .

Theorem 6.4. Under Assumptions 3 and 5,  $E(Z|H_0) = \sqrt{2 \log P H_n}$ .

*Theorem 6.5.* Under  $n \to \infty$  and  $H_0$ , given the properties of  $S_0$  and  $S_1$ , then:

$$S|H_0 \rightarrow^d N(0,1),$$

the power of *S* is approaching 1 once the arbitrage characteristic is selected as:

$$\Pr(\operatorname{reject} H_0 | \mathcal{M} \neq \emptyset) \to 1.$$

#### 7. Numerical Study

In this section, we use Compustat and Fama-French three and five factors data to simulate stock returns and then evaluate the performance of our estimation and hypothesis test procedures.

#### 7.1. Data Generation

First, we use Fama-French three factors monthly returns and all the characteristics that will be included in the empirical study to simulate the stock excess returns. Most of the characteristics are updated annually so we treat those variables as time-invariant during each one-year rolling block. For the characteristics that are updated every month, we substitute the mean values as their fixed values for each fiscal year. We use Fama-French monthly returns from July of year *t* to June of year t+1 and characteristics of fiscal year t-1 to generate the stock returns from July of year *t* to June of year t + 1. The periods we generate are the same

Table 1. Simulation Results 1 Part 1.

		Distortion%	%0	%0	%0	%0	%0	35.9%	%0	%0	0.1%	%0	1.3%	%0	%0	%0	%0	0.2%	%0	%0	%0	%0	%0	%0	%0	%0	%0	%0	%0	%0	%0	0.3%	0.2%	%0	%0	0.2%	%0	%0	%0	0.1%	%0	%0	
		%	%(	%(	3%	2%	3%	1%	%€	%(	%(	7%	1%	3.7	1%	%(	4%	%	%6	%(	%(	5%	%(	%	3%	%(	%(	%(	%(	%(	%(	%(	%(	%(	%(	%(	3%	3%	%0	%€	1%	%(	
		Selected			7.		82.	68.	99.	10	10	.66	98.	9	57.	10	0.5	68.	97.	10	10	92.	10	÷.	27.	10	10	10	10	10	10	10	10	10	2	10	10	10	10	.66	1.1%	-	
		MSE	8.1	3.7	13.7	-6.2	4.1	17.5	8.4	5.4	5.6	5.7	7.8	4.1	3.7	6.6	10.8	20	11	6.4	7.4	5.4	4.2	4.2	5.5	3.2	4.9	5.4	3.4	5.5	5.5	8.5	7.8	7.4	11.2	23.3	9.4	3.6	7.1	4.4	4	4.4	
		S1	-8.2	-8.5	1.7-	0.06	-7.2	5.4	4.7	-5.8	'n	-5.3	0	-7.5	-7.5	-5.3	6.8	-4.9	-6.2	4.3	-6.7	-5.2	-6.4	-6.1	6.3	-5.4	-5.7	4.5	4.1	4.	6.6	7.8	6.7	-1.2	-1.	1.7	1.9	1.4	-a.1	-2.3	-7.6	-8.1	
		So	0	0	0.7		7.4	10.9	6	10	6	8.9	8.9	5.7	5.1	6	4.8	6.1	8.8	11	1	10.2	11	0.1	m	=	11	11	11	11	1	1	11	11	1	1	11	11	11	11	0.1	0	
	Mode	S	-8.2	-8.5	<i>L</i> -	-6.1	0.2	16.3	4.3	4.2	9	3.6	8.9	-1.8	-2.4	3.7	-7	1.2	2.6	6.7	4.3	ŝ	4.6	ę	-3.3	5.6	5.3	6.5	6.9	6.9	17.6	18.8	17.7	9.8	9.7	12.7	12.9	12.4	7.9	8.7	-7.5	-8.1	
	Semiparametric Mode	Distortion%	87.4%	2.1%	%0	%0	6.5%	73.4%	0.02%	6.9%	32.4%	61.1%	12.9%	7.1%	15.3%	2.3%	0.6%	%0	8.4%	NA	20.3%	%0	28.2%	3.5%	7.1%	%0	%0	NA	NA	%0	%0	%0	%0	48.2%	%0	%0	%0	%0	%0	0.3%	0.3%	%0	
$^{2} = 4$	Sei	elected % D	100%	100%	100%	%0	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	NA	100%	100%	100%	100%	100%	100%	%0	NA	NA	%0	100%	2.5%	100%	100%	100%	34.4%	87.3%	4.5%	100%	100%	100%	100%	
ď		MSE 9	8.6	4.3	14.1	12.5	4.5	8.4	8.6	5.5	5.8	5.9	8.3	4.6	4	6.9	1.1	20.6	11.5	ΝA	∞	6.1	4.9	Ŝ	6.3	4	5.8	ΝA	ΝA	6.5	6.4	9.3	8.6	8.1	1.9	24	10.1	4.4	7.9	5.3	5.3	5.2	
		S <sub>1</sub> N																																							2.3		
	del	So	10.8																																								
	inear Mode	S																																							9.1		
	Lin	rtion%																																							%0		
		Distor																																									
	del	Selected %	81.2%	99.9%	100%	96.8%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	
	iparametric Model	MSE	9	1.6	11.5	9.5	7	15.8	6.2	3.2	3.5	3.6	5.8	7	1.6	4.5	8.7	17.9	6	4.3	5.2	3.2	7	2.1	з.3	-	2.7	3.3	1.2	3.3	3.4	6.3	5.7	5.2	8.9	21.2	7.2	1.4	4.9	2.3	1.7	2.2	
	aramet	S1																																							0		
	Semipi	So	6.2	∞	6	9.1	6	12.6	6	10	6	6	9.1	6	6	6	6	6	~	11	_	_	11	11.6	11	=	11	12.9	11			~	_	11.3		~ .		11	11.2	11.3	11	11	
		S	-0.5	3.4	8.6	9.2	10.6	28.6	19.5	20.3	23.1	16.2	20.7	10.6	10.6	15.8	11.5	9.9	13.2	23.2	17.8	22.4	20.2	17.2	16.3	23.1	20.6	24.9	27.3	26.2	58.2	59.2	61	33.2	42.7	37.3	41.3	41.3	26.5	28.9	1	9.5	
$\sigma^2 = 1$		Distortion%	100%	100%	%0	%0	100%	100%	74.2%	100%	100%	100%	85.1%	100%	100%	100%	30.4%	%0	100%	NA	100%	%0	100%	99.5%	54.2%	69.9%	98.6%	NA	NA	%0	%0	%0	%0	100%	%0	%0	%0	%0	%0	66.8%	0.2%	%0	
	del	Selected % [	100%	100%	100%	%0	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	NA	100%	100%	100%	100%	100%	100%	66.7%	NA	NA	%0	100%	80.4%	100%	100%	100%	99.7%	100%	100%	100%	100%	100%	100%	
	Linear Mod	MSE Se	4.0	2	ە	.2	31	16	4.9		9.6	2.7	.1	38	73	17	6.9	5.4	2.0	Ą	8.	8.	7	78	5	ø.	5.5	٩A	٩A	5	2	5	4.	6.9	9.6	22	6.		9.0	m	2.9	6.9	
	Line	S <sub>1</sub> M	Ι.																_		-	-								_				-			-	_					
																																									7 16.1		
		s S	9 11.5		_			_											_						_	_	_			_			_			_							
			24.	32.	34.	-0.5	9	б	78.	13.	117	90.	84.	83.	113.	88.	45	20.	59.	Ż	56.	43	59.	40	41.	<u>-</u>	15.	Ż	Ż	0	10.	<u>.</u>	<u>1</u> .	23.	.6	9	80	10.	14.	19.	17	21.	
		2	468	894	1108	1199	1333	1409	1466	1560	1494	1292	1393	1340	1285	1181	1110	1044	1125	2192	2236	2273	2235	2270	2405	2376	2323	2344	2434	2548	2741	2928	2894	2905	2804	2570	2516	2491	2402	2326	2241	2178	
		Window	-	2	m	4	Ŝ	9	7	8	6	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	

			ī	Linear model			settibal attrettic titodet			-	i	Linear model	ē			Semiparametric model	tric moc	e				
'indow n 41 2113	5 24.1 é	50	51 M 18	ISE Selected % 1.7 100%	dow n S 5 <sub>0</sub> 5 <sub>1</sub> MSE Selected % Distortion% 41 2113 24.1 6.1 18 4.7 100% 7.5%	5 7.8	50 10	51 N -2.2	MSE Se 3.9	Selected % Distortion% 100% 0%	Distortion% 0%	5 50 13.9 5.7	50 51 5.7 8.2	51 MSE .2 6.9		Selected % Distortion% 100% 0%	6 S 6 -8.1	5 50 1 0	-8.1	MSE Se	E Selected % 0%	Selected % Distortion% 0% 0%
	18.4 5	5.7 12			و 0%	11.3	10	1.3	9	100%	%0	10.8	5.8 5	1.	100%		Ľ	1.0				%0
2007	18.8 5	5.7 13			و 0%	9.1	10	-0.9	4.1	100%	%0	10.5	5.7 4				6 -8.3	m	8			%0
1924		5.8 10				13.6	10.8	2.8	7.5	100%	8%	11.2	5.8 5	-				5 2.	φ			0.2%
1990	27.5 5	5.7 21				8.1	10	-1.9	1.4	100%	%0	13.3	577					8	~ ~			%0
46 1937	20.3 5	5.8 14	14.5 5	5.4 100%	6 0.9%	19.7	11.8	7.9	4.7	100%	18%	12.6 5	5.9 6.	6.7 7.6		6 3%		8 11.	2 -3.2	6.8	100%	12.3%
1909	13.2 5	5.7 7				14.2	10.4	3.8	4.5	100%	3.5%	8.8	5.7 3				6 2.7	7 8.	4			%0
1872	21.8 5	5.7 16				11.4	10	1.4	2	100%	%0	11.1	5.8 5					8	5 -7.			%0
1841	16.3 5	5.7 10	5.0	2.1 100%		8.7	10	-1.3	1.4	100%	0.1%	8.1	5.7 2	4.4			6 -8.4	4	~; ~			%0
	11 5	5.7 5		4.3 100%	و 0%	12.6	10.6	2	3.5	100%	3.5%	6.5	5.7 0	.8 6.6	5 99.7%	0		6	φ ο			%0

Table 2. Simulation Results 1 Part 2.

as the empirical study, namely, 50 years from July 1967 to June 2017. For each rolling block with 12 months, we have

$$y_{it} = h(X_i) + \sum_{j=1}^{3} g_j(X_{ij}) f_{jt} + \epsilon_{it},$$
 (7)

 $y_{it}$  is the generated stock's return;  $h(X_i)$  is the mispricing function consisting of a nonlinear characteristic function of  $x_i$ , which is to mimic the sparse structure of the mispricing function;  $g_j(X_{ij})$  is the *j*th characteristics-based factor loading, which has an additive semiparametric structure;  $X_{ij}$  is the *j*th subset consisting of 4 characteristics;  $f_{jt}$  is the *j*th Fama-French factor returns at time t;  $\epsilon_{it}$  is the idiosyncratic shock for stock *i* at time t, generated from  $N(0, \sigma^2)$ .

We generate characteristic functions

$$h(X_i) = \sin X_i,$$

$$g_j(\mathbf{X}_{ij}) = X_{ij1}^2 + (3X_{ij2}^3 - 2X_{ij2}^2) + (3X_{ij3}^3 - 2X_{ij3}) + X_{ij4}^2,$$

where  $X_{ijl}$  is a randomly picked characteristic without replacement from the data in empirical study and j = 1, 2, 3, l = 1, ..., 4. A description of these characteristics can be found in the appendix. Additionally, all  $h(X_i)$ ,  $g_j(X_{ij})$  are adjusted to have zero mean and unit variance. As we use real data to conduct the simulation, the Assumption of independent X may not be satisfied. Although some characteristics are correlated, the semiparametric model overcomes this problem properly when compared with the parametric model that has serious size distortion.

We do not specify  $h(X_i)$  and  $g_j(X_{ij})$  to be orthogonal explicitly, but we draw characteristics without replacement and employ sine-waves and polynomials to approximate the orthogonality as much as possible. In this simulation, our method can only estimate the component of  $h(X_i)$  that is orthogonal to  $g_j(X_{ij})$ . However, results reveal that one can still select the arbitrage characteristics even if we cannot estimate arbitrary  $h(X_i)$  unrestrictively.

#### 7.2. Model Misspecification

In this subsection, we show the necessity to consider semiparametric analysis when the forms of factor loadings and mispricing functions are nonlinear.

Under the data generation process, we consider both semiparametric and linear analysis to compare Mean Squared Error (MSE) and hypothesis test results under both specifications. We apply our estimation methodology in Section 3 to estimate Equation 7. For semiparametric specification, we choose the number of B-splines bases to be  $\lfloor n^{0.3} \rfloor$ . *n* is the number of assets in each balanced rolling window, and  $\lfloor \cdot \rceil$  means the nearest integer. We orthogonalize these bases, and then use the Projected-PCA and restricted OLS to estimate model Equation (7). As for the hypothesis test part, we choose threshold value to be  $\eta_n = H_n \sqrt{2 \log(PH_n)} = \lfloor n^{0.3} \rceil \sqrt{2 \log(P\lfloor n^{0.3} \rceil)}$ , where *P* is the number of characteristics, and *n* is the number of stock in each rolling block. For the linear specification, each characteristic only has one basis, which is itself. In terms of the hypothesis test,

<u></u> :
÷
F
Pa
5
÷
Res
æ
L
0
Ξ
-10
2
⊢
S
m'
Ð
-
He he
E.

		88	%	%	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	%	%	%	%	%	%	%	%	%	%	%(	%	%	0/1	%	%	3%	%	%	%(	%	%	%	%	%	%	%	%	%	%	%	%	%
		Distortion% 0%	0%	 -	-	10		100	<u></u>	0			0	-	0	01	<u> </u>	2	5	101	2.2	94.8	0	<u>0</u>	<u>ě</u>	01	0	0	-		<u>p</u>	<u>0</u>	<u>10</u>	-			<u>0</u>	10
		Selected % 0%	12.9%	73.5%	100 %	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	%001	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	13.3%
		MSE Se 6.9	1 1 1 2		- 4	. 7	1.7	6.7	4.6	t 8	2.7	2.5	9	0.2	8.0	ю. С	0,7	- 0	j o	0 -	- <del>.</del> .	2.7	3.9	6.4 1.0	1.7	4 /	0.0	2.5	0.5	9.4	2.3	80.00	0.8	 	6.1	4.0	2.4	3.4
		S <sub>1</sub> N -8.5	-7.5							_																			-					_				
	۳ ا	مى	c	יי. מית					18																													
	factors Ĵ =	-8.5 -8.5	-6.5 5 0	0 c																																		
	ted fac		ī			10		2	~ ~	1						21	0 7 0	- 4	D	4		2		4	ο I	Ω,	43	(	7		m I	LO I	ŝ	-	9	4 (	2	2
= 5	Number of estimated	Distortion% 1.3%	%0 0%	%C.U	%0.c.z	%06	0.3%	0.5%	1.4%	17.2%	%0	0.1%	0.2%	0.1%	2.1%	7.1%	%9.//	%0 702 0	0/ 70	%C.0 26.7%	%0	%0	%0	54%	2.6%	0.2%	36.8%	52.2%	45.9%	15.7%	85.6%	28.9%	0.4%	3%	29.2%	9.4%	%0	1.3%
Number of factors J =	Numb	Selected % 97.4%	100%	001	%001 %001	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	%001 100%	%001 70001	10006	100%	100%	100%	100%	100%	100%	%0	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%
umber	5 :	MSE 6	2.3	0.0	200	13.5	9.2	Ś	4.4	4 10 0 9	1.8	1.4	4.7	9.8	17.8	10.7	0 r 4 •	0 4 4. c	4 c	7 K	1 m 1 m	1.2	2.8	3.2	m.	m i	τ. 	4. 1	5.5	6.1	10.1	41.2	18.4	1.6	5.1	2.8	1.7	2.2
z	ors $\hat{J} =$	5 <sub>1</sub> -2.4	-2.5	0.0	9	30.4	17.3	20.4	22.9	19.3	6.2	6.1	12.4	8.1	5.7	14.4	41.7	- 6	19.4	01 0	10.7	19.3	15.8	19.1	27	23.8	4 7	65.9	60.9	28.9	33.6	34.9	27.4	39.2	23.2	55	3.9	1.8
	ed fact	5 <sub>0</sub> 6.9	∞ ⊂	11 א	6	28.3	6	10	9.2	10.7	6	6	6	6	9.2	9.7	78.0	= :		1100	51	1	1	17	11.3	=;	15.4	18./	16.3	12.8	20.4	14.5	=	11.3	14.2	12.2	Ξ	11.2
	stimat	5.4.5	5.5	14.5	167	58.7	26.3	30.4	32.1	30.5	15.2	15.1	21.4	17.1	14.9	24.1	69.8 1	1.02	0.00	43.7	21.7	30.3	26.8	36.1	38.3	34.8	/9.4	84.6	11.2	41.7	54.1	49.4	38.4	50.5	37.4	37.2	14.9	12.9
	Number of estimated factors $\hat{J}$	Distortion% 0%	%0 %0	0.00	%0 *	28%	%0	%0	%0 0	1.1%	%0	%0	%0	%0	%0	%0 0%	%0	%0	040	%0 %0	%0	%0	%0	17.1%	%0	%0 7	1.3%	/.8%	21.6%	3%	68.5%	10.4%	0.4%	0.4%	2.2%	2.1%	%0	0.3%
	ors $\hat{J}=3$	Selected % 81.2%	99.9%	06 8%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%
	estimated factors.	MSE 6	1.6	U N	j c	15.8	6.2	3.2		0.0	7	1.6	4.5	8.7	17.9	9,	4 r v c	7.0	4 C	7 r		-	2.7	3.3	1.2	m v	4.5	0.7	5.7	5.2	8.9	21.2	7.2	1.4	4.9	2.3	1.7	2.2
	stimate	5 <sub>1</sub> -5.7	-4.6	- - -	- 9	15.9	10.5	10.3	14.1	11.6	1.6	1.6	6.8	2.5	0.9	4.2	7.7	0.8	t. c	7 Y Y	5 G	12.1	9.6	12	16.3	15.2	47.1	47.4	47.6	21.9	24.2	25.1	30.3	30.3	15.3	17.6	0	-1.5
	Ψ	5 <sub>0</sub> 6.2	∞ ⊂	ם ۲ ע	- o	12.6	6	10	<b>о</b> с	9 ر 1 ر	6	6	6	6	6	, ہ	= ;	= =	= =	11 -	2	11	1	12.9	=	; ;		8. II 8. II	13.4	11.3	18.5	12.2	1	1	11.2	11.3	: 3	=
۳ ا	Number of	-0.5	3.4	0.0	10.6	28.6	19.5	20.3	23.1	20.7	10.6	10.6	15.8	11.5	6.6	13.2	72.7	8.7	1. C	20.2	16.3	23.1	20.6	24.9	27.3	26.2	58.2	59.2	61	33.2	42.7	37.3	41.3	41.3	26.5	28.9	; ;	9.5
Number of factors J	= 5	Distortion% 0%	%0 %1	0.1.0 0%	01%	51.3%	0.01%	%0	0.7%	0.1% 4%	%0	0.2%	0.2%	%0	0.8%	0.1%	0.7% 0.2%	%0 0 407	0.4.%	0%0 23 q%	0.1%	0.1%	0.4%	51.3%	0.9%	0.1%	10.9%	%0	45.4%	16%	86.8%	%0	2.9%	6.8%	10.5%	10.3%	0.1%	1.1%
Num	estimated factors $\hat{J}$	Selected % 99.6%	99.9% 10001	00 80%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	0001	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%	24.1%	100%	100%	100%	27.2%	100%	100%	100%	100%	100%	100%
	f estim.	MSE 5.9	1.5	4. O	, c	15.6	6.1	m	с. С. с	5. T	1.8	1.4	4.4	8.5	17.8	8. v	4 L	- °	n 0	<u>o</u> 0	3.2	0.8	2.5	3.1	-	2.7 M	3.2	0.1	5.5	Ś	8.8	21.1	2	1.3	4.7	2.1	1.6	7
	Number of	-4.4	2° -	0 C	44	24.1	17.8	17.6	22.6	18.4	6.2	6.4	12.9	7.4	4.9	9.7	20.8	10.4	10.4 16.5	C.01	11.5	19.6	16.1	19.7	25.2	23.5	60.7	60.1	61	29.5	33.3	33.4	39.6	39.5	22.2	25.1	3.8	7
	Nun	50 7	∞ ⊂	10 ع د 10 ع	0	17.5	6	10	9.1	9 م 4	6	6	6	6	9.1	ه 5	= ;	= :	= =	137	<u>;</u> =	11	11.1	16.7	11.1	13	12.3	13.8	16.3	12.9	20.5	14.2	11.3	11.8	12.2	12.3	=	11.1
		S 2.6	1 6 1	12.0	1.4	41.6	26.8	27.6	31.7	C.22	15.2	15.4	21.9	16.4	13.3	18.7	x . x	24.4 7 0 0 0	4. CZ	070	22.5	30.6	27.2	36.4	36.3	34.5	2	/3.9	11.3	42.4	53.8	47.6	50.9	51.3	34.4	37.4	14.8	13.1
		468 n	894	1100	1333	1409	1466	1560	1494	1393	1340	1285	1181	1110	1044	1125	7617	2230	2000	0202	2405	2376	2323	2344	2434	2548	2/41	2928	2894	2905	2804	2570	2516	2491	2402	2326	2241	2178
		Window 1	2 0	04	t vr	9	7	8	6	2 =	12	13	14	15	16	11	<u>8</u> 2	910	02 50	17	23	24	25	26	27	28	57	0	 	32	ŝ	34	35	36	37	38	39	40

art 2.	
mulation Results 2 Pa	
Table 4. Sir	

Number of factors J = 3

Number of factors J = 5

			ΠN	5		Number of estimated factors J = 5	= 5	Number of estimated factors $J = 3$	5		מומרו			Number of estimated factors $J = 5$	ed ract	ors J =	<u>.</u>	INUME	Number of estimated factors $J = 1$	1 factors	/ = 3				
Vindow 41	dow n S S <sub>0</sub> 41 2113 11 10	2 E	50 10	5 <u>1</u>	MSE Se 3.8	S1         MSE         Selected %         Distortion%           1         3.8         100%         0.2%	Distortion% 0.2%	S 7.8	100°	51 1 -2.2	MSE S 3.9	Selected % 100%	Distortion% 0%	5 11.5	10°5	5.1 1.5	MSE 4.5	Selected % 100%	Distortion% 0.2%	5 50 41.1 32.4	5 <sub>0</sub> 32.4	5 <sub>1</sub> 8.7	MSE 5.4	Selected % 99.9%	Distortion% 100%
42		15.2	10	5.2	5.9	100%	%0	11.3	10	1.3	9	100%	%0	15.7	10	5.7	6.4	100%	%0	φ	0	ø	9.2	%0	60
43	2007 1	12.6	10	2.6	4	100%	0.5%	9.1	10	-0.9	4.1	100%	%0	13.4	10.2	3.2	4.7	100%	1.7%	-0.1	6.4	-6.5	5.6	64.4%	%0
4	1924 1	1 6.91	13.1	6.8	7.3	100%	30.7%	13.6	10.8	2.8	7.5	100%	8%	19.5	12.9	6.6	7.5	100%	28.9%	20	20	0	8.3	100%	100%
45	1990 1	11.4	10	1.4	1.2	100%	0.1%	8.1		-1.9	1.4	100%	%0	20.7	14.6	9	1.8	100%	45.2%	116	20	96	1.7	100%	1009
46	1937 2	27.1	14	13.1	4.5	100%	37.7%	19.7		7.9	4.7	100%	18%	28.3	14.8	13.5	5.4	100%	45.8%	24.6	20	4.6	6.2	100%	1009
47		19.5 1	11.7	7.8	4.4	100%	16.1%	14.2		3.8	4.5	100%	3.5%	24	14	10	4.4	100%	38.1%	51.7	35.2	16.5	5.4	100%	1009
48		15.2	10	5.1	1.8	100%	0.2%	11.4		1.4	7	100%	%0	15	10	S	2.1	100%	0.1%	5	10	ņ	2.9	100%	60
49		12.3 1	10.1	2.2	1.2	100%	1.1%	8.7		-1.3	1.4	100%	0.1%	11.8	10.1	1.7	4.4	100%	0.8%	-10	0	-10	4.4	%0	60
50		18.5 1	12.4	6.1	3.3	100%	15%	12.6	10.6	2	3.5	100%	3.5%	20.2	13.3	6.9	3.7	100%	19.3%	-3.9	0.4	4.3	4.4	3.9%	0

we use the same logic as in the semiparametric settings, and we set  $\eta_n = \sqrt{3 \log(P)}$ .

In all the estimation above, we assume that we know the real number of factors, which is three. We will discuss the situation when the number of factors is unknown in the next subsection. Mean Squared Error (MSE) is also reported to compare the fitness of models (Equation (7)).

From Table 1, under different noise levels, namely  $\sigma^2 = 1$  and  $\sigma^2 = 4$ , the semiparametric model outperforms the linear model in the following aspects:

- 1. The fitness of the semiparametric model is much better than the linear model, which can be illustrated from MSE.
- 2. The semiparametric model can enhance the power of  $S_1$  by non-zero  $S_0$ , which can not only select the correct mispricing characteristics but also avoid size distortions. As for the linear model, it is influenced by the correlated characteristics. Therefore, during certain periods we even obtain the noninvertible characteristic matrix. The linear model can also select the relevant covariates with decent probability, but it suffers from serious size distortions. In contrast, our semiparametric model with orthogonal bases can mitigate this problem to a great extent.
- 3. Because  $S_1$  can be very small and even negative, especially when the noise  $\sigma_i$  is strong, the additional component  $S_0$ is necessary to strengthen the power of  $S_1$  and to select the relevant characteristics that can explain the mispricing function.

#### 7.3. Robustness Under Stronger Noise

In Table 1, we set two different noise levels of random shocks, namely  $\sigma^2 = 1$  and  $\sigma^2 = 4$ . Although  $\sigma^2 = 1$  is closer to the empirical data, we conduct this comparison to show the robustness of our methods. When the noise level becomes three times bigger, the accuracy of power-enhanced tests gets much lower for certain windows. However, there are no size distortions under comparatively high noise level recalling that all the components of our simulation model are rescaled to have unit variance. Another fact is that the stronger noise does deteriorate the power of conventional Wald tests, leading to an even smaller value of  $S_1$ , which can be mitigated through adding  $S_0$ .

Therefore, we conclude that our methods are robust to a higher noise level regarding no size distortions. However, the accuracy of selecting relevant components and the role of enhancing the power of hypothesis tests will be influenced negatively.

#### 7.4. Number of Factors

In the empirical study, the number of factors is unknown. Therefore, in this subsection, we will study whether our methodology is robust to a various numbers of factors considered.

We simulate according to another data-generation process

$$y_{it} = h(X_i) + \sum_{j=1}^{5} g_j(X_{ij}) f_{jt} + \epsilon_{it},$$
 (8)

similarly,  $y_{it}$  is the generated stock return;  $h(X_i)$  is the mispricing function consisting of a nonlinear characteristic function of  $X_i$ , to mimic the sparse structure of the mispricing function;  $g_j(X_{ij})$  is the *j*th characteristics-based factor loading, which has an additive semiparametric structure;  $X_{ij}$  is a subset consisting of four characteristics;  $f_{jt}$  is the *j* Fama-French 5-factor returns at time t;  $\epsilon_{it}$  is the idiosyncratic shock, generated from  $N(0, \sigma^2)$ . Moreover, we generate characteristic functions as follows:

$$h(X_i) = \sin X_i$$

$$g_j(\mathbf{X}_{ij}) = X_{ij1}^2 + (3X_{ij2}^3 - 2X_{ij2}^2) + (3X_{ij3}^3 - 2X_{ij3}) + X_{ij4}^2$$

where  $X_{ijl}$  is a randomly picked characteristic without replacement from the data in empirical study with j = 1, ..., 5, l = 1, ..., 4. Furthermore, all  $h(X_i)$  and  $g_j(X_{ij})$  are adjusted to have zero mean and unit variance.

Given the above data-generation process, we test the influence of over and under-estimated number of factors. We choose the number of factors to be either three or five, and compare the results in Table 3. The first category column is the scenario of over estimating the number of factors. We simulate the data generation process using the Fama-French three factors model, but estimate the number of factors to be five. However, this does not cause any serious problems. For some rolling blocks, the probability of mistakenly selected irrelevant characteristics is slightly higher under over estimating the number of factors. Moreover, over estimating the number of factors can increase the model fitting marginally. Therefore, we conclude that over estimating the number of factors does not cause severe size distortion using our methods.

On the other hand, under estimating the number of factors can lead to misleading test results. We can conclude this from the last several columns where we estimate the number of factors to be three in a five-factor model. Compared with the correct specified model, under estimating causes not only higher MSE, but also higher distortions, which means it is more likely to select irrelevant characteristics. Therefore, in the empirical study we prefer the five-factor model rather than the three-factor model.

#### 8. Empirical Study

#### 8.1. Data

We use monthly stock returns from CRSP and firms' characteristics from Compustat, ranged from 1965 to 2017. We construct 33 characteristics following the methods of Freyberger et al. (2020). Details of these characteristics can be found in the Appendix. We use characteristics from fiscal year t - 1 to explain stock returns between July of year t to June of year t + 1. After adjusting the dates from the balance sheet data, we merge two datasets from CRSP and Compustat. We require all firms included in our analysis to have at least three years of characteristics data in Compustat.

Data is modified with regards to the following aspects:

1. Delisting is quite common for CRSP data. We use the way of Hou et al. (2015) to correct the returns of all delisted assets before 2018. Detailed methods can be found in their appendix.

- 2. Projected-PCA works well, even under small *T* circumstances. Thus, we choose the width of our window to be 12 months. Another reason for the short window width is that we assume that mispricing functions are time-invariant in each window. One of the limitations of Projected-PCA is that it can only be used for a balanced panel, which means the number of stock will vary when we applied one-year rolling windows to obtain a short-time balanced panel. Meanwhile, we take monthly updated characteristics' mean values of 12 months as fixed characteristic values in each window. We also use the rolling window method to detect peer groups of arbitrage characteristics.
- 3. B-splines are based on each time-invariant characteristic of *n* firms, which are not delisted in each window.
- Rolling windows are moving at a 12-month step from July 1967 to June 2017 without overlapping. The first 24-month returns are not included as they do not have corresponding characteristics.
- 5. Excess returns are obtained by the difference between monthly stock returns and monthly Fama-French risk-free returns.

#### 8.2. Estimation

We construct B-splines bases based on evenly distributed knots, and the degree of each basis is three. We choose  $H_n = \lfloor n^{0.3} \rfloor$ , and *n* is the number of stocks. To get a relatively large balanced panel in each window, some characteristics with too many missing values are eliminated. Therefore, only 33 characteristics are left. Firms kept in balanced panels in our dataset range from 468 to 2928, which means that both *n* and  $\hat{A} \in \mathbb{R}^{PH_n}$  are diverging. Large *n* can satisfy asymptotic requirements. These facts emphasize the necessity of introducing a power-enhanced component into the hypothesis test. Before the next step, we use time-demeaning matrix  $D_T$  to demean excess return matrix in each window.

Next, we project the time-demeaned monthly excess return matrix  $\tilde{Y}$  onto the B-splines space spanned by characteristics bases  $\Phi(X)$ , and then we collect the fitted values  $\hat{Y}$ . We apply principal component analysis on  $\frac{1}{n}\hat{Y}^{\mathsf{T}}\hat{Y}$ , and attain the first five eigenvectors corresponding to the first five biggest eigenvalues as the estimates of unobservable factors F. We choose the number of factors to be five according to simulation results.

Then, we estimate the factor loading matrix by:

$$\hat{G}(X) = \hat{Y}\hat{F}(\hat{F}^{\mathsf{T}}\hat{F})^{-1}$$

Moreover, we use equality-constrained OLS to estimate the mispricing function. We project excess monthly return matrix on the characteristic space  $\Phi(X)$  that is orthogonal to factor loading matrix  $\hat{G}(X)$ .

Another goal of this article is to conduct a power-enhanced test on the mispricing function. Therefore, our final step is to estimate the covariance matrix  $\Sigma$  of  $\hat{A}$ .

#### 8.3. Power-Enhanced Hypothesis Tests

In this section, we conduct a power-enhanced test in each rolling block. First, we set threshold value for each window,  $\eta_n =$ 

 $\lfloor n^{0.2} \rceil \sqrt{2 \log(PH_n)}$ , where  $H_n$  is the number of bases for each characteristic whereas *P* is the number of total characteristics in each window, with P = 33.  $\eta_n$  is data-driven critical value and it diverges as the number of firms increases. We use indicator function  $I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n)$  to achieve three goals.

- 1. This indicator function selects the most relevant characteristics that can explain the variation of the mispricing function. Results of the last column in Table 5 are characteristics selected in  $\hat{\mathcal{M}} = \{X_p \in X : \sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n, p = 1, 2, \ldots, p\}.$
- 2. It contributes to the test statistic *S* by adding a diverging power-enhanced component  $S_0$ . As T = 12 is small in this empirical study, we assume the homoscedasticity of  $\epsilon_{it} + \gamma_i$ . We also specify an over-shrunk covariance matrix by setting off-diagonal elements to be zeros.
- 3. It avoids size-distortion by the conservative critical value  $\eta_n$ .

The diagonal elements of  $\hat{\Sigma}$  are estimated variances of mispricing coefficients. These elements can be substituted into the indicator function  $I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n)$ , where  $\hat{\sigma}_{ph}$  is the *ph*th diagonal element of  $\hat{\Sigma}$ .

Finally, the new statistic *S* can be calculated as follows:

$$S = S_0 + S_1,$$

$$S_0 = H_n \sum_{p=1}^{P} I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \ge \eta_n), \quad S_1 = \frac{\hat{A}^{\mathsf{T}} \hat{\Sigma}^{-1} \hat{A} - PH_n}{\sqrt{2PH_n}}.$$

#### 8.4. Test Results

This section presents the empirical results. Details can be found in Table 5, which lists the results of 50 rolling windows from Jul.1967 to Jun.2017. Generally, the number of firms included in the 12-month rolling block is increasing period by period. The number of our characteristic B-splines bases is a function of the number of firms *n* in each block. Therefore, the dimension of the mispricing coefficient vector  $\mathbf{A} \in \mathbb{R}^{PH_n}$  is also diverging. This verifies the necessity of using a power-enhanced component  $S_0$ .

Recalling that  $S|H_0 \rightarrow {}^d N(0, 1)$ , some of the test statistics S is big enough to reject the null hypothesis. However, for some testing windows, there are no strong signals showing the existence of characteristics-based mispricing functions after subtracting systematic effects. Most  $S_1$  values are small and even negative, which may be caused by the sparsity structure of the mispricing function or/and the low power problems due to diverging dimension of mispricing coefficients.

The power-enhanced component  $S_0$  works well in the empirical study. It selects the most important explanatory characteristics and strengthens the power of  $S_1$ , mitigating the low power problem.

Apart from contributing to the power of tests, the indicator function in the power-enhanced component can also screen out the most relevant characteristics, which are concluded as "Characteristics Selected" in Table 5.

Some empirical findings are worth discussing. Although short-term cumulative returns like  $r_{2_1}$  are always selected, we cannot take this as the evidence of arbitrage opportunities since we construct  $r_{2_1}$  as the time-invariant average of all  $r_{2_1}$  in

Table 5. Empirical study results.

Time period	n	S	S <sub>0</sub>	S <sub>1</sub>	MSE	Characteristics selected
Jul.1967–Jun.1968	468	-9.6	0	-9.6	0.005	NONE
Jul.1968–Jun.1969	951	-0.45	8	-8.45	0.004	r <sub>2_1</sub>
Jul.1969–Jun.1970	1108	1.7	9	-7.3	0.005	r <sub>2</sub> 1
Jul.1970–Jun.1971	1199	-8.7	0	-8.7	0.006	NONE
Jul.1971–Jun.1972	1333	-10	0	-10	0.004	NONE
Jul.1972–Jun.1973	1409	12.7	18	-5.3	0.005	r <sub>12</sub> 2,r <sub>6</sub> 2
Jul.1973–Jun.1974	1466	2.1	9	-6.9	0.005	r <sub>2</sub> 1
Jul.1974–Jun.1975	1560	-10.7	0	-10.7	0.01	NONE
Jul.1975–Jun.1976	1494	0.1	9	8.9	0.05	<i>r</i> <sub>2 1</sub>
Jul.1976–Jun.1977	1292	0.1	9	-9	0.004	r <sub>2</sub> 1
Jul.1977–Jun.1978	1393	-9.4	0	-9.4	0.005	NONE
Jul.1978–Jun.1979	1340	8.6	18	-9.4	0.005	r2_1,r12_7
Jul.1979–Jun.1980	1285	1	9	-8	0.005	r2_1/ 12_/
Jul.1980–Jun.1981	1181	9.7	18	-8.2	0.006	r <sub>12</sub> 7,r <sub>12</sub> 2
Jul.1981–Jun.1982	1110	1.2	9	-7.8	0.01	r2_// r2_2
Jul.1982–Jun.1983	1044	33.1	36	-3	0.01	r12_2,r12_7,r6_2,r2_1
Jul.1983–Jun.1984	1125	-0.9	9	-9.9	0.006	r2_1
Jul.1984–Jun.1985	2192	-0.2	11	-11.2	0.01	r <sub>2</sub> 1
Jul.1985–Jun.1986	2236	13.1	22	-8.94	0.01	r <sub>12</sub> 7,r <sub>12</sub> 2
Jul.1986–Jun.1987	2273	1.7	11	-9.3	0.01	PCM
Jul.1987–Jun.1988	2235	0.9	11	-10.1	0.01	r <sub>2</sub> 1
Jul.1988–Jun.1989	2270	1.2	11	-9.8	0.01	RÔĂ
Jul.1989–Jun.1990	2405	-0.1	11	-11.1	0.01	r <sub>2_1</sub>
Jul.1990–Jun.1991	2376	1.1	11	-9.9	0.02	r <sub>2</sub> 1
Jul.1991–Jun.1992	2323	2.1	11	-8.9	0.02	r <sub>2_1</sub>
Jul.1992–Jun.1993	2344	12.2	22	-9.8	0.02	r <sub>12</sub> 7,r <sub>12</sub> 2
Jul.1993–Jun.1994	2434	0.4	11	-10.6	0.01	r <sub>2</sub> _7, r <sub>2_2</sub>
Jul.1994–Jun.1995	2548	2.4	11	-8.6	0.01	r2_1
Jul.1995–Jun.1996	2741	14.1	22	-7.9	0.02	BEME,r <sub>2_1</sub>
Jul.1996–Jun.1997	2928	18.1	22	-3.9	0.01	BEME, r <sub>2</sub>
Jul.1997–Jun.1998	2894	26.5	33	-6.5	0.02	r2 1,r12 7,r12 2
Jul.1998–Jun.1999	2905	24.6	33	-8.4	0.02	AT,LME,r <sub>2_1</sub>
Jul.1999–Jun.2000	2804	13.8	22	-8.2	0.03	r <sub>2</sub> 1,r <sub>12</sub> 7
Jul.2000–Jun.2001	2570	37.7	44	-6.3	0.02	AT,LME, r <sub>2_1</sub> , r <sub>6_2</sub>
Jul.2001–Jun.2002	2516	1.3	11	-9.7	0.02	r2_1
Jul.2002–Jun.2003	2491	15	22	-7	0.02	Lev, $r_{2}^{2}$
Jul.2003–Jun.2004	2402	3.9	11	-7.1	0.01	r <sub>2</sub> 1
Jul.2004–Jun.2005	2326	1.8	11	-9.2	0.01	IPM
Jul.2005–Jun.2006	2241	2.5	11	-8.5	0.01	r <sub>2_1</sub>
Jul.2006–Jun.2007	2178	1.5	11	-9.5	0.01	r <sub>2 1</sub>
Jul.2007–Jun.2008	2113	12.6	20	-7.4	0.01	r <sub>12_2</sub> ,r <sub>2_1</sub>
Jul.2008–Jun.2009	2023	1.7	10	-8.3	0.02	r2_2, 2_1
Jul.2009–Jun.2010	2007	1	10	-9	0.01	r2_1
Jul.2010–Jun.2011	1924	13.6	20	-6.4	0.01	r2_1
Jul.2011–Jun.2012	1990	2.5	10	-7.5	0.01	r <sub>2_1</sub>
Jul.2012–Jun.2013	1937	23.7	30	-6.3	0.01	r2_1,r12_7,r12_2
Jul.2013–Jun.2014	1909	2.3	10	-7.7	0.01	r2_1//12_7//12_2
Jul.2014–Jun.2015	1872	5.5	10	-4.5	0.01	r <sub>2_1</sub>
Jul.2015–Jun.2016	1841	12.4	20	-7.6	0.01	DelGmSale,r <sub>2_1</sub>
Jul.2016–Jun.2017	1826	26.1	30	-3.9	0.01	C2D,PCM,r <sub>12_7</sub>
						· · · · · · · · · · · · · · · · · ·

NOTE: This table summarizes the empirical results, where n represents the number of stock in this rolling window.

the same rolling window, which contains much overlapping information of monthly excess returns. However, this is not the case for long-term and mid-term cumulative returns like  $r_{12_2}$ ,  $r_{12_7}$  and  $r_{6_2}$ , because these average returns include a lot of information from another rolling window (Figure A.1).

Apart from the cumulative returns, some other characteristics contribute to the arbitrage opportunities as well. PCM (price to cost margin) appears twice. From Figure A.2, we find that the PCM mispricing curve is nonlinear and generally decreasing as the value of PCM increases. ROA (Return-On-Asset) also plays a role during 1988–1989. It behaves like a parabola with fluctuations near zero in Figure A.3. As for Lev (ratio of longterm debt and debt in the current liabilities), it is decreasing for Lev 0 and increasing afterward as in Figure A.7. In Figure A.8, IPM (pretax profit margin) function behaves like a "V" shape with the turning point zero during 2004–2005. DelGmSale (difference in the percentage in Gross margin and the percentage change in Sales) experiences a bump at zero during 2015– 2016 in Figure A.9. C2D (cash flow to price) curve behaves like "V" around the zero in 2016-2017, (see Figure A.10). All characteristics curves in the above figures are standardized as uniformly distributed characteristics in the interval [-100, 100]. This is for presentation purposes only since most characteristics are unevenly distributed.

Another finding is the persistence of some arbitrage characteristics. Arbitrage characteristics can be persistent for two years once appeared, such as BEME (ratio of the book value of equity and market value of equity) in Figure A.4. Some persistent arbitrage characteristics even have similar shapes of mispricing functions in different rolling windows, such as AT (Total asset) in Figure A.6 and LME (total market capitalization of the previous month) in Figure A.5.

#### 8.5. Dynamic Peer Groups of Arbitrage Characteristics


In this section, we illustrate that there are distinguishable peer groups of the same arbitrage characteristic resulting in similar mispricing returns. We apply the methods in Section 5 and take two rolling windows, namely, Jul.1986–Jun.1987 and Jul.2004–Jun.2005 as demonstrative examples.

In the rolling window July1986–June 1987, PCM is selected as the only arbitrage characteristic that can explain arbitrage returns. We reveal that similar characteristic-based arbitrage returns are determined by distinguishable groups of the characteristic PCM. We first divide arbitrage returns  $\ddot{y}_{it}$  into different return groups. And then, we detect whether there are some clustering structures within groups of the highest and the lowest characteristic-based arbitrage returns, respectively. As we have 2326 assets, for the visualization purpose, we set the threshold value of the K-means method to be relatively small to have as many as ten groups.

In Table 6, Group 2 has the largest positive average return while Group 6 has the worst. Next, we detect the clusterings of characteristic "PCM" within each group individually, which is the second layer in Section 5.

In Table 7, there are two clusterings of PCM that provide the highest positive characteristic-based arbitrage returns. Group 2.2, which has an extreme negative PCM value but a high characteristic-based arbitrage return, is an outlier. Members in group 2.1 with excellent arbitrage performance have positive and small PCM values.

Table 8 gives groups of PCM in group 6. Members of this group are divided into two clusterings. Group 6.1 has a relatively large PCM value, while Group 6.2 has a smaller PCM, which is close to that in group 2.1 with the highest arbitrage return. This is an evident illustration of the nonlinear structure of h(X) in this window. The structure of characteristic-based arbitrage returns during July 1986– June 1987 is:




The classification can be found at Figure A.11, where assets are labeled by their "PERMNO," and both axes are rescaled.

Another example is the characteristic-based arbitrage return  $\ddot{y}_{it}$  during the year 2004–2005. The power-enhanced test selects characteristic "IPM" as the only explanatory variable.

We apply the Hierarchical K-means method. The results of the first layer classification can be found in Table 9. There are ten groups in total according to the similarity of characteristicbased arbitrage returns. Next, we pick two groups with the highest and the lowest returns, respectively, to give clusterings of "IPM" in these two groups.

Similarly, we show classification results in Tables 10 and 11 . Positive IPM values give higher characteristic-based arbitrage returns. On the contrary, when IPM is close to zero or negative, the characteristic-based arbitrage returns fall into the lowest group (Group 8).



The plots of the IPM can be found at Figure A.12, where the axes are rescaled, and assets are labeled by their "PERMNO" code with five digits.

Finally, it is obvious that peer groups of arbitrage characteristics are dynamic in two aspects. First, the selected arbitrage characteristics are time-varying. Although some of the arbitrage characteristics can show up for more than one block once they appeared, no arbitrage characteristic can be substantially persistent. Second, as in Figure A.4, the same arbitrage characteristic can have different functional forms in various rolling windows. However, the patterns of some characteristics show persistence in different time periods, such as AT in Figure A.6 and LME in Figure A.5. In a word, under the flexible semiparametric setting, methods for constructing arbitrage portfolio in Kim et al. (2021) are inapplicable, although the characteristics-based mispricing function is significant for certain time periods.

**Table 6.** First layer 1986–1987 (clusterings of  $\ddot{y}_{it}$ ).

Group number	Group centroid	Group size
1	0.0059	435
2	0.1205	26
3	-0.0082	428
4	0.0399	189
5	0.0697	71
6	-0.1018	29
7	-0.0617	110
8	-0.0390	250
9	-0.0225	349
10	0.0208	386

Table 7.	Second layer	1986-1987	(clusterings of characteris	tic
PCM).				

Group	Centroids of Arbitrage returns	Centroids	Group
number		of PCM	size
2.1	0.1211	0.2452	25
2.2	0.1039	—7.630	1

Table 8. Second layer 1986-1987 (clusterings of characte	ristic
PCM).	

Group	Centroids of Arbitrage returns	Centroids	Group
number		of PCM	size
6.1	-0.1085	0.728	9
6.2	-0.0989	0.288	20

**Table 9.** First layer 2004–2005 (clusterings of  $\ddot{y}_{it}$ ).

Group centroid	Group size
0.0421	276
0.0059	459
0.1537	26
-0.024	367
0.0659	166
0.023	387
0.0999	120
-0.0758	67
-0.0437	244
-0.0082	436
	0.0421 0.0059 0.1537 -0.024 0.0659 0.023 0.0999 -0.0758 -0.0437

 Table 10.
 Second layer 2004–2005 (clusterings of characteristic IPM)

Group	Centroids of Arbitrage	Centroids	Group
number	returns	of PCM	size
3.1	0.1681	0.266	5
3.2	0.1502	0.143	21

Table 11. Second layer 2004–2005 (clusterings of characteristic IPM).

Group number	Centroids of arbitrage returns	Centroids of PCM	Group size
8.1	-0.0713	-0.07	10
8.2	-0.1016	-0.134	57

#### 9. Conclusion


We proposed a semiparametric characteristics-based factor asset pricing model, with a focus on the existence and the structure of the mispricing function. Both unknown characteristicsbased factor loadings and the mispricing component are approximated by B-splines sieve. The model is estimated by both Project-PCA and equality-constrained OLS. We also develop a power-enhanced test to investigate whether there are mispricing characteristics, orthogonal to the main systematic factors. This is necessary because when the B-splines coefficients of the mispricing function are diverging, the conventional Wald test has very low power. The traditional Wald test is strengthened by a screening process, which adds significant components to the original statistic. Our proposed methods work well for both simulations and the U.S. stock market. Empirically, we find distinct clusterings of the same characteristics resulting in similar arbitrage returns, forming "peer groups." The mispricing functions are time-varying and mostly insignificant under our setting.

## Appendix

### A1. Characteristic Description

#### Table A.1. Characteristic details.

Name	Description	Reference
A2ME	We define assets-market cap as total assets (AT) over market capitalization as of December t-1. Market capitalization is the product of shares outstanding (SHROUT) and price(PRC).	Bhandari (1988)
AT ATO	Total assets (AT) Net sales over lagged net operating assets. Net operating assets are the difference between operating assets and operating liabilities. Operating assets are total assets (AT) minus cash and short-term investments (CHE), minus investment and other advances (IVAO). Operating liabilities are total assets (AT), minus debt in current liabilities(DLC),minus long-term debt (DLTT),minus minority interest (MIB), minus preferred stock (PSTK), minus common equity (CEQ).	Gandhi and Lusting (2015) Soliman(2008)
BEME	Ratio of book value of equity to market value of equity. Book equity is shareholder equity (SH) plus deferred taxes and investment tax credit (TXDITC), minus preferred stock (PS). SH is shareholder's equity (SEQ). If missing, SH is the sum of common equity (CEQ) and preferred stock (PS). If missing, SH is the difference between total assets (AT) and total liabilities (LT). Depending on availability, we use the redemption (item PSTKRV), liquidating (item PSTKL), or par value (item PSTK) for PS. The market value of equity is as of December t-1. The market value of equity is the product of shares outstanding (SHROUT) and price (PRC).	Rosenberg, Reid and Lanstein (1985) Davis, Fama, and French (2000)
2D	Ratio of cash and short-term investments (CHE) to total assets (AT) Cash flow to price is the ratio of income and extraoridinary items (IB) and depreciation and amortization (dp) to total liabilities (LT).	
CTO Debt2P	We define caoital turnover as ratio of net sales (SALE) to lagged total assets (AT). Debt to price is the radio of long-term debt (DLTT) and debt in current liabilities (DLC) to the market capitalization as of December t-1. Market capitalization is the product of shares outstanding (SHROUT) and price (PRC).	Haugen and Baker (1996) Litzenberger and Ramaswamy (1979)
$\Delta ceq$ $\Delta(\Delta Gm - Sales)$	The percentage change in the book value of equity (CEQ). The difference in the percentage change in gross margin and the percentage change in sales (SALE). We define gross margin as the difference in sales (SALE) and costs of goods sold (COGS).	Richardson et al. (2005) Abarbanell and Bushee (1997)
∖Shrout ∖PI2A	The definition of the percentage change in shares outstanding (SHROUT). We define the change in property, plants ,and equipment as changes in property,plants,and equipment (PPEGT) and inventory (INVT) over lagged total assets (TA).	Pontiff and Woodgate (2008) Lyandres , Sun, and Zhang (2008)
DTO	We define turnover as ratio of daily volume (VOL) to shares outstanding (SHROUT) minus the daily market turnover and de-trend it by its 180 trading day median. We scale down the volume of NASDAQ securities by 38% after 1997 and by 50% before that to address the issue of double-counting of volume for NASDAQ securities.	Garfinkel (2009); Anderson and Dyl (2005)
2P	We define earnings to price as the ratio of income before extraordinary items (IB) to the market capitalization as December t-1 Market capitalization is the product of share outstanding (SHROUT) and price (PRC).	Basu (1983)
EPS	We define earnings per share as the ratio of income before extraordinary items (IB) to share outstanding (SHROUT) as of December t-1	Basu (1997)
nvestment PM	We define investment as the percentage year-on-year growth rate in total assets (AT). We define pretax profit margin as ratio of pretax income (PI) to sales (SALE).	Cooper, Gulen and Schill(2008)
ev	leverage is the ratio of long-term debt (DLTT) and debt in the current liabilities (DLC) to the sum of long-term debt, debt in current liabilities, and stockholders' equity (SEQ)	Lewenllen (2015)
.ME	Size is the total market capitalization of the previous month defined as price (PRC) times shares outstanding (SHROUT)	Fama and French (1992)
Furnover PCM	Turnover is last month's volume (VOL) over shares outstanding (SHROUT). The price-to-cost margin is the difference between net sales (SALE) and costs of goods sold (COGS) divided by net sales (SALE).	Datar, Naik and Radcliffe (1998) Gorodnichenko and Weber (2016) and D'Acunto, Liu, Pflucger and Wcber (2017)
M Q	The profit margin is operating income after depreciation (OIADP) over sales (SALE) Tobin's Q is total assets (AT), the market value of equity (SHROUT times PRC) minus cash and short-term investments (CEQ) minus deferred taxes (TXDB) scaled by total assets (AT).	Soliman (2008)
ROA ROC	Return-on-assets is income before extraordinary items (IB) to lagged total assets (AT). ROC is the ratio of market value of equity (ME) plus long-term debt (DLTT)minus total assets to Cash and Short-Term Investments (CHE).	Balakrishnan, Bartov and Faurel (2010) Chandrashekar and Rao (2009)
ROE 12-2	Return-on-equity is income before extraordinary items (IB) to lagged book-value of equity. We define momentum as cumulative return from 12 months before the return prediction to two months before.	in Haugen and Baker (1996) Fama and French (1996)
12-7	We define intermediate momentum as cumulative return from 12 months before the return prediction to seven months before.	Novy-Marx (2012)
6–2	We define $f_{6-2}$ as cumulative return from 6 months before the return prediction to two months before.	Jegadeesh and Titman (1993)
2-1	We define short-term reversal as lagged one-month return.	Jegadeesh(1990)
S2C	Sales-to-cash is the ratio of net sales (SALE) to Cash and Short-Term Investments (CHE).	following Ou and Penman (1989)
Sales-G SGA2S	Sales growth is the percentage growth rate in annual sales (SALE). SGA to sales is the ratio of selling ,general and administrative expenses (XSGA) to net sales	Lakonishok, Shleifer , and Vishmy (1994
JUNZJ	(SALE).	



0.4

-0.6

0 PCM

(b) PCM Curve 2016-2017



0.10

0 PCM

(a) PCM Curve 1984-1985

-50






Figure A.3. Mispricing characteristic curve of standardized ROA in 1988–1989.



(a) BEME Curve 1995-1996



(b) BEME Curve 1996-1997

Figure A.4. Mispricing characteristic curve of standardized BEME.




Figure A.5. Mispricing characteristic curve of standardized LME.




Figure A.6. Mispricing characteristic curve of standardized AT.




Figure A.7. Mispricing characteristic curve of standardized LEV in 2002–2003.




Figure A.8. Mispricing characteristic curve of standardized IPM in 2004–2005.




Figure A.9. Mispricing characteristic curve of standardized DelGmSale in 2015–2016.






Figure A.10. Mispricing characteristic curve of standardized C2D in 2016–2017.



(a) Clustering of PCM with highest returns

(b) Clustering of PCM with lowest returns

Figure A.11. Clustering of PCM 1986–1987



(a) Clustering of IPM with highest returns

Figure A.12. Clustering of IPM 2004–2005.

#### A3. Proofs

Throughout the proofs, we have the number of observations  $n \to \infty$ , and time T is fixed.

Proof of Theorem 6.1. In Equation (5), we have

$$Y = (\Phi(X)A + \Gamma + R^{\mu}(X))\mathbf{1}_{T}^{\mathsf{T}} + (\Phi(X)B + \Lambda + R^{\theta}(X))F^{\mathsf{T}} + U,$$

Multiply time-demeaned matrix  $D_T$  on both sides, where  $D_T = I_T$  –  $\frac{1}{T} \mathbf{1}_T^{\mathsf{T}} \mathbf{1}_T$ . Given time-invariant mispricing components, we obtain:

$$YD_T = (\Phi(X)B + \Lambda + R^{\theta}(X))F^{\mathsf{T}}D_T + UD_T.$$

On-wards, we define  $YD_T = \tilde{Y}$  and  $F^{\mathsf{T}} = F^{\mathsf{T}}D_T$ . Time-demeaned factors do not change their properties.

Next, multiple both sides by  $P = \Phi(X)(\Phi(X)^{\mathsf{T}}\Phi(X))^{-1}\Phi(X)^{\mathsf{T}}$ ,

$$\ddot{Y} = (\Phi(X)B + P\Lambda + PR^{\theta}(X))F^{\mathsf{T}} + PUD_T.$$

We decompose

 $P\tilde{Y} = \hat{Y} = \Phi(X)BF^{\mathsf{T}} + P\Lambda F^{\mathsf{T}} + PUD_T + PR^{\theta}(X)F^{\mathsf{T}} = e_1 + e_2 + e_3 + e_4,$ 

as  $n \to \infty$  and  $n^{\nu} \to \infty$ , approximation error  $\mathbf{R}^{\boldsymbol{\theta}}(\mathbf{X}) \to_{P} \mathbf{0}$  as in Huang et al. (2010). Thus,  $\mathbf{e}_{\mathbf{1}}^{\mathsf{T}} \to^{P} \mathbf{0}$ . Under Assumption 1, we have following results: for  $\frac{1}{n} \sum_{j=1}^{3} \mathbf{e}_{\mathbf{2}}^{\mathsf{T}} \mathbf{e}_{j}$ ,


$$\frac{1}{n}\boldsymbol{P}\Lambda \rightarrow^{\boldsymbol{P}} \boldsymbol{0},$$

therefore,

$$\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{2}^{\mathsf{T}}}\boldsymbol{e_{j}}+\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{j}^{\mathsf{T}}}\boldsymbol{e_{2}}\rightarrow^{P}\boldsymbol{0}$$

For  $\frac{1}{n} \sum_{j=1}^{3} e_{3}^{\mathsf{T}} e_{j}$ ,

$$\frac{1}{n}PU \rightarrow^{P} \mathbf{0},$$



(b) Clustering of IPM with lowest returns

therefore,

$$\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{2}^{\mathsf{T}}}\boldsymbol{e_{j}}+\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{j}^{\mathsf{T}}}\boldsymbol{e_{2}}\rightarrow^{P}\boldsymbol{0}.$$

And only  $\frac{1}{n}e_1^{\mathsf{T}}e_1$  left, namely

$$\frac{1}{n}e_1^{\mathsf{T}}e_1 = F\frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n}F^{\mathsf{T}}.$$

Under Assumption 2–4 and fixed T. A much smaller  $T \times T$ matrix  $\frac{1}{n}\hat{Y}^{\mathsf{T}}\hat{Y}$  can be sovied by asymptotic principal component by Connor and Korajczyk (1986).  $\hat{F} = \frac{1}{\sqrt{T}} \{\psi_1, \psi_2, \dots, \psi_J\},\$ where  $\{\psi_1, \psi_2, \dots, \psi_J\}$  are eigenvectors corresponding to the first J eigenvalues of  $\frac{1}{n} \hat{Y}^{\mathsf{T}} \hat{Y}$ .

Thus,  $\hat{F} \rightarrow P F$  follows.  $\Box$ 

*Proof of Theorem 6.2.* Given  $\hat{F}$ , we have

$$\hat{\boldsymbol{G}}(\boldsymbol{X}) = \hat{\boldsymbol{Y}}\hat{\boldsymbol{F}}(\hat{\boldsymbol{F}}^{\mathsf{T}}\hat{\boldsymbol{F}})^{-1}$$

as  $\hat{F}^{\mathsf{T}}\hat{F} = I_I$ , therefore,

$$\hat{G}(X) = \tilde{Y}\hat{F}$$

Then we need to show

$$\mathbb{E}((\hat{G}(X_i) - G(X_i))^2) = 0.$$

Take the sample analogue

$$\frac{1}{n}(\hat{G}(X) - G(X))^{\mathsf{T}}(\hat{G}(X) - G(X)).$$

Given

$$G(X) = \Phi(X)B + R^{\theta}(X).$$

$$\hat{G}(X) = (\Phi(X)B + P\Lambda + PR^{\theta}(X))F^{\mathsf{T}}\hat{F} + PUD_T\hat{F}$$

Furthermore,

$$G(X) - \hat{G}(X) = (\Phi(X)B + P\Lambda + PR^{\theta}(X))F^{\mathsf{T}}\hat{F} + PUD_T\hat{F}$$
$$- \Phi(X)B - R^{\theta}(X) = q_1 + q_2 + q_3 + q_4.$$

Similar to the Proof of Theorem 6.1,

$$\frac{1}{n}(\hat{G}(X) - G(X))^{\mathsf{T}}(\hat{G}(X) - G(X)) \to^{P} \frac{1}{n}q_{1}^{\mathsf{T}}q_{1}$$
$$+ \frac{1}{n}q_{3}^{\mathsf{T}}q_{3} + \frac{1}{n}q_{1}^{\mathsf{T}}q_{3} + \frac{1}{n}q_{3}^{\mathsf{T}}q_{1}.$$

For the first term,

$$\frac{1}{n}q_1^{\mathsf{T}}q_1 = \hat{F}^{\mathsf{T}}F(\Phi(X)B + P\Lambda + PR^{\theta}(X))^{\mathsf{T}}(\Phi(X)B + P\Lambda + PR^{\theta}(X))F^{\mathsf{T}}\hat{F}$$

due to

$$\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{2}^{\mathsf{T}}}\boldsymbol{e_{j}}+\frac{1}{n}\sum_{j=1}^{3}\boldsymbol{e_{j}^{\mathsf{T}}}\boldsymbol{e_{2}}\rightarrow^{P}\boldsymbol{0},$$

and

 $\frac{1}{n}e_1^Te_1 \to^P F \frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n}F^{\mathsf{T}}$ 

then,

$$\frac{1}{n}q_1^Tq_1 \to^P \hat{F}^{\mathsf{T}}F\frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n}F^{\mathsf{T}}\hat{F}.$$

Theorem 6.1 and Assumption 2 give  $\hat{F} \to F$  and  $F^T F = I_J$ , therefore

$$\frac{1}{n}q_1^Tq_1 \rightarrow^P \frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n},$$

Similarly,

$$\frac{1}{n}q_3^Tq_3 \to^P \frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n},$$
$$\frac{1}{n}q_1^Tq_3 \to^P -\frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n},$$
$$\frac{1}{n}q_3^Tq_1 \to^P -\frac{B^{\mathsf{T}}\Phi^{\mathsf{T}}(X)\Phi(X)B}{n}.$$

Therefore,

$$\frac{1}{n}q_{1}^{\mathsf{T}}q_{1} + \frac{1}{n}q_{3}^{\mathsf{T}}q_{3} + \frac{1}{n}q_{1}^{\mathsf{T}}q_{3} + \frac{1}{n}q_{3}^{\mathsf{T}}q_{1} \to 0.$$

Then,

 $\frac{1}{n}(\hat{G}(X) - G(X))^{\mathsf{T}}(\hat{G}(X) - G(X)) \to^{P} 0,$ 

thus,

$$\hat{G}(X) \rightarrow^{P} G(X).$$

Then Theorem 6.2 follows.

*Proof of Theorem 6.3.* Let  $\dot{Y} = \frac{1}{T}(Y - \hat{G}(X)\hat{F})\mathbf{1}_T$ . By substituting the restriction, we have the Lagrangian equation:

$$\min_{A} (\dot{Y} - \Phi(X)A)^{\mathsf{T}} (\dot{Y} - \Phi(X)A) + \lambda \hat{G}^{\mathsf{T}}(X)\Phi(X)A \qquad (A.1)$$

Then we take the first order condition with respect to A and  $\lambda$  separately, and we obtain

$$\begin{pmatrix} 2\Phi(X)^{\mathsf{T}}\Phi(X) & \Phi(X)^{\mathsf{T}}\Theta(X) \\ \hat{G}(X)^{\mathsf{T}}\Phi(X)^{\mathsf{T}} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \hat{A} \\ \lambda \end{pmatrix} = \begin{pmatrix} 2\Phi(X)^{\mathsf{T}}\Phi(X)^{\mathsf{T}}\Phi(X) \\ \mathbf{0} \end{pmatrix}.$$
(A.2)

Under Assumption 2, the above matrices are invertible, which can be written as follows:

$$\begin{pmatrix} \hat{A} \\ \lambda \end{pmatrix} = \begin{pmatrix} 2\Phi(\mathbf{X})^{\mathsf{T}}\Phi(\mathbf{X}) & \Phi(\mathbf{X})^{\mathsf{T}}\mathbf{Q}(\mathbf{X}) \\ \hat{G}(\mathbf{X})^{\mathsf{T}}\Phi(\mathbf{X})^{\mathsf{T}} & \mathbf{0} \end{pmatrix}^{-1} \begin{pmatrix} 2\Phi(\mathbf{X})^{\mathsf{T}}\mathbf{P} \\ \mathbf{0} \end{pmatrix}.$$
(A.3)

Therefore, we obtain

$$\hat{A} = Q\tilde{A},$$

where

$$\boldsymbol{Q} = \boldsymbol{I} - (\boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}} \boldsymbol{\Phi}(\boldsymbol{X}))^{-1} \boldsymbol{\Phi}(\boldsymbol{X})^{\mathsf{T}} \hat{\boldsymbol{G}}(\boldsymbol{X}) (\hat{\boldsymbol{G}}(\boldsymbol{X})^{\mathsf{T}} \hat{\boldsymbol{G}}(\boldsymbol{X}))^{-1} \hat{\boldsymbol{G}}(\boldsymbol{X})^{\mathsf{T}} \boldsymbol{\Phi}(\boldsymbol{X}),$$

$$\tilde{A} = \frac{1}{T} (\Phi(X)^{\mathsf{T}} \Phi(X))^{-1} \Phi(X)^{\mathsf{T}} \dot{Y} \mathbf{1}_{T}.$$

Furthermore, let  $\Xi = \Phi(\mathbf{X})\mathbf{A} - h(\mathbf{X}) = \Phi(\mathbf{X})\mathbf{Q}\mathbf{A} - \Phi(\mathbf{X})$  $\mathbf{A} - \mathbf{R}^{\mu}(\mathbf{X}).$ 

Under the restriction  $\hat{G}(X)^{\mathsf{T}} \Phi(X) A = 0$ , we can obtain

$$\Xi = \Phi(\mathbf{X})\mathbf{M}(\Phi(\mathbf{X})^{\mathsf{T}}\Phi(\mathbf{X}))^{-1}\Phi(\mathbf{X})^{\mathsf{T}}\frac{1}{T}(\Phi(\mathbf{X})\mathbf{A} + \mathbf{R}^{\mu}(\mathbf{X}) + \Gamma + (\Lambda + \mathbf{R}^{\theta}(\mathbf{X}))\mathbf{F}')\mathbf{1}_{T} - \Phi(\mathbf{X})\mathbf{A} - \mathbf{R}^{\mu}(\mathbf{X}).$$
(A.4)

Furthermore, we have:

$$\Phi(\mathbf{X})\mathbf{M}(\Phi(\mathbf{X})^{\mathsf{T}}\Phi(\mathbf{X}))^{-1}\Phi(\mathbf{X})^{\mathsf{T}} = (\mathbf{I} - \Phi(\mathbf{X})(\Phi(\mathbf{X})^{\mathsf{T}}\Phi(\mathbf{X}))^{-1}$$
$$\Phi(\mathbf{X})^{\mathsf{T}}\hat{\mathbf{G}}(\mathbf{X})(\hat{\mathbf{G}}(\mathbf{X})^{\mathsf{T}}\hat{\mathbf{G}}(\mathbf{X}))^{-1}$$
$$\hat{\mathbf{G}}(\mathbf{X})^{\mathsf{T}})\mathbf{P}. \tag{A.5}$$

And then, substitute Equation A.5 into Equation A.4 and under Assumption 1 and Theorem 6.2

$$\Xi = \Phi(\mathbf{X})\mathbf{A} - \Phi(\mathbf{X})\mathbf{A} - \mathbf{R}^{\mu}(\mathbf{X}).$$

$$R^{\mu}(X) \to \mathbf{0}$$
 as  $n \to \infty$ ,

therefore,

$$\frac{1}{n} \Xi^{\mathsf{T}} \Xi \to \mathbf{0}$$

And the Theorem 6.3 follows.  $\Box$ 

Proof of Theorem 6.4.

Define 
$$Z = \max_{\{1 \le p \le P, 1 \le h \le H_n\}} \{|\hat{\alpha}_{ph}| / \hat{\sigma}_{ph}\}$$
. Under Assumption 3, we have

$$\hat{\alpha}_{ph}/\hat{\sigma}_{ph}|H_0 \rightarrow^d N(0,1).$$

Therefore, under the  $H_0$ , we have:  $\rho^{tE(Z)} < F[\rho^{tZ_1}]$ 

$$= E[max\{t|\hat{\alpha}_{ph}|/\hat{\sigma}_{ph}\}]$$

$$= E[max\{t|\hat{\alpha}_{ph}|/\hat{\sigma}_{ph}\}]$$

$$= P_{h,h=1} E[e^{t|\hat{\alpha}_{ph}|/\hat{\sigma}_{ph}}]$$

$$= PH_n e^{t^2/2}.$$

Then take the logarithm of both sides we can obtain

$$E[Z] \leqslant \frac{\log PH_n}{t} + \frac{t}{2}.$$

If we set 
$$t = \sqrt{2 \log PH_n}$$
 to minimise  $\frac{\log PH_n}{t} + \frac{t}{2}$ , then we have:

$$E[Z] \leqslant \sqrt{2 \log PH_n}.$$

Therefore, we can bound the  $|\hat{\alpha}_{ph}|/\hat{\sigma}_{ph}$  by  $\sqrt{2 \log PH_n}$ . *Proof of Theorem 6.5.* To proof

$$\Pr(\operatorname{reject} H_0 | \hat{\mathcal{M}} \neq \emptyset) \to 1,$$

equivalently, we need to prove

$$\Pr(S_0 + S_1 | \hat{\mathcal{M}} \neq \emptyset) \to 1$$

$$\begin{split} S_0 &= H_n \sum_{p=1}^P I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \geqslant \eta_n), \text{ as } H_n = n^{\nu} \to \infty \text{ when } \\ n \to \infty. \\ \text{Once } \hat{\mathcal{M}} \neq \emptyset, \text{ then } \sum_{p=1}^P I(\sum_{h=1}^{H_n} |\hat{\alpha}_{ph}| / \hat{\sigma}_{ph} \geqslant \eta_n) \geqslant 1, \text{ therefore, } \\ S_0 \to \infty \text{ as } n \to \infty. \text{ Meanwhile } F_q = O(1), \text{ we can show that:} \end{split}$$

$$\Pr(S_0 + S_1 > F_q | \hat{\mathcal{M}} \neq \emptyset) \to 1.$$

Then the Theorem 6.5 follows.  $\Box$ 

#### **Supplementary Materials**

The supplement material includes a table of characteristics description, proofs, plots of mispricing curves, and figures of peer groups.

#### **Acknowledgments**

Shuyi Ge thanks the Fundamental Research Funds for the Central Universities for research support; Shaoran Li thanks the School of Economics, Peking University for research support.

#### References

- Carhart, M. M. (1997), "On Persistence in Mutual Fund Performance," *The Journal of Finance*, 52, 57–82. [1]
- Chen, X., and Pouzo, D. (2012), "Estimation of Nonparametric Conditional Moment Models With Possibly Nonsmooth Generalized Residuals," *Econometrica*, 80, 277–321. [3]
- Connor, G., Hagmann, M., and Linton, O. (2012), "Efficient Semiparametric Estimation of the Fama–French Model and Extensions," *Econometrica*, 80, 713–754. [2,3,5]
- Connor, G., and Korajczyk, R. A. (1986), "Performance Measurement With the Arbitrage Pricing Theory: A New Framework for Analysis," *Journal* of Financial Economics, 15, 373–394. [22]
- Connor, G., and Linton, O. (2007), "Semiparametric Estimation of a Characteristic-Based Factor Model of Common Stock Returns," *Journal* of Empirical Finance, 14, 694–717. [2]
- Cox, D. R. (1957), "Note on Grouping," Journal of the American Statistical Association, 52, 543–547. [6]
- Fama, E. F., and French, K. R. (1993), "Common Risk Factors in the Returns on Stocks and Bonds," *Journal of Financial Economics*, 33, 3–56. [1,5]

(2015), "A Five-Factor Asset Pricing Model," Journal of Financial Economics, 116, 1–22. [1,5]

- Fan, J., Liao, Y., and Mincheva, M. (2013), "Large Covariance Estimation by Thresholding Principal Orthogonal Complements," *Journal of the Royal Statistical Society*, Series B, 75, 603–680. [5]
- Fan, J., Liao, Y., and Wang, W. (2016), "Projected Principal Component Analysis in Factor Models," *Annals of Statistics*, 44, 219. [2,3,4,8]
- Fan, J., Liao, Y., and Yao, J. (2015), "Power Enhancement in High-Dimensional Cross-Sectional Tests," *Econometrica*, 83, 1497–1541. [1,2,5,6]

- Feng, G., Giglio, S., and Xiu, D. (2020), "Taming the Factor Zoo: A Test of New Factors," *The Journal of Finance*, 75, 1327–1370. [1]
- Fisher, W. D. (1958), 'On Grouping for Maximum Homogeneity," Journal of the American Statistical Association, 53, 789–798. [6]
- Freyberger, J., Neuhierl, A., and Weber, M. (2020), "Dissecting Characteristics Nonparametrically," *The Review of Financial Studies*, 33, Pages 2326 -2377. [1,2,13]
- Hjalmarsson, E., and Manchev, P. (2012), "Characteristic-Based Mean-Variance Portfolio Choice," *Journal of Banking & Finance*, 36, 1392–1401. [1,2,6]
- Hoberg, G., and Phillips, G. (2016), "Text-Based Network Industries and Endogenous Product Differentiation," *Journal of Political Economy*, 124, 1423–1465. [2]
- Hou, K., Xue, C., and Zhang, L. (2015), "Digesting Anomalies: An Investment Approach," *The Review of Financial Studies*, 28, 650–705. [1,13]
- Huang, J., Horowitz, J. L., and Wei, F. (2010), "Variable Selection in Nonparametric Additive Models," *Annals of Statistics*, 38, 2282. [2,3,4,22]
- Kelly, B. T., Pruitt, S., and Su, Y. (2017), "Instrumented Principal Component Analysis," Available at SSRN 2983919. [2]
- (2019), "Characteristics Are Covariances: A Unified Model of Risk and Return," *Journal of Financial Economics*, 134, 501–524. [2,5]
- Kim, S., Korajczyk, R. A., and Neuhierl, A. (2021), "Arbitrage Portfolios," *The Review of Financial Studies*, 34, 2813–2856. [1,2,4,5,6,16]
- Kock, A. B., and Preinerstorfer, D. (2019), "Power in High-Dimensional Testing Problems," *Econometrica*, 87, 1055–1069. [2,5]
- Ledoit, O., and Wolf, M. (2012), "Nonlinear Shrinkage Estimation of Large-Dimensional Covariance Matrices," *The Annals of Statistics*, 40, 1024– 1060. [5]
- Liew, C. K. (1976), "Inequality Constrained Least-Squares Estimation," Journal of the American Statistical Association, 71, 746–751. [5]
- Pesaran, M. H., and Yamagata, T. (2012), "Testing CAPM With a Large Number of Assets," in AFA 2013 San Diego Meetings Paper. [5]
- Pollard, D. (1981), "Strong Consistency of k-Means Clustering," *The Annals of Statistics*, 9, 135–140. [7]
- (1982), "A Central Limit Theorem for k-Means Clustering," The Annals of Probability, 10, 919–926. [7]
- Ross, S. (1976), "The Arbitrage Theory of Capital Asset Pricing," Journal of Economic Theory, 13, 341–360. [1]
- Sun, W., Wang, J., and Fang, Y. (2012), "Regularized k-Means Clustering of High-Dimensional Data and Its Asymptotic Consistency," *Electronic Journal of Statistics*, 6, 148–167. [7]
- Vogt, M., and Linton, O. (2017), "Classification of Non-Parametric Regression Functions in Longitudinal Data Models," *Journal of the Royal Statistical Society*, Series B, 79, 5–27. [3,7]